

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



ML335

Item No. 27635

CAS Registry No.:	825658-06-8	
Formal Name:	N-[(2,4-dichlorophenyl)methyl]-4-	CI
	[(methylsulfonyl)amino]-benzamide	0
MF:	C ₁₅ H ₁₄ Cl ₂ N ₂ O ₃ S	
FW:	373.3	
Purity:	≥98%	S I I I I
UV/Vis.:	λ _{max} : 258 nm	
Supplied as:	A solid	l H
Storage:	-20°C	
Stability:	≥2 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis		

Laboratory Procedures

ML335 is supplied as a solid. A stock solution may be made by dissolving the ML335 in the solvent of choice, which should be purged with an inert gas. ML335 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of ML335 in ethanol is approximately 5 mg/ml and approximately 30 mg/ml in DMSO and DMF.

ML335 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, ML335 should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. ML335 has a solubility of approximately 0.04 mg/ml in a 1:20 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

ML335 is an activator of the two-pore domain potassium channels $K_{2p}2.1/TREK1$ and $K_{2p}10.1/TREK2$ (EC₅₀s = 14.3 and 5.2 μ M, respectively, in *Xenopus* oocytes).¹ It is selective for $K_{2p}2.1/TREK1$ and $K_{2p}10.1/TREK2$ channels over $K_{2p}4.1/TRAAK$ channels. ML335 activates K_{2P}^{-2} 2.1/TREK1 by binding to the C-type gate, which is the active site of TREK channels.

Reference

1. Lolicato, M., Arrigoni, C., Mori, T., et al. K_{2P}2.1(TREK-1): Activator complexes reveal a cryptic selectivity filter binding site. Nature 547(7663), 364-368 (2017).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

SAFETY DATA This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/23/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM