

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com


PRODUCT INFORMATION



Pentadecanoic Acid-d₂₀

Item No. 28081

CAS Registry No.:	130348-95-7
Formal Name:	pentadecanoic-d ₂₉ acid
Synonyms:	C15:0-d ₂₉ , Pentadecylic Acid-d ₂₉
MF:	$C_{15}HD_{29}O_{2}$
FW:	271.6
Chemical Purity:	≥95% (Pentadecanoic Acid)
Deuterium	
Incorporation:	≥99% deuterated forms (d_1 - d_{29}); ≤1% d_0
Supplied as:	A solid
Storage:	-20°C
Stability:	≥2 years
Information represents the product specifications. Batch specific analytical resu	



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Pentadecanoic acid-d₂₉ is intended for use as an internal standard for the quantification of pentadecanoic acid (Item No. 17399) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Pentadecanoic acid-d₂₉ is supplied as a solid. A stock solution may be made by dissolving the pentadecanoic acid-d₂₉ in the solvent of choice. Pentadecanoic acid-d₂₉ is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of pentadecanoic acid-d₂₀ in ethanol and DMF is approximately 25 mg/ml and approximately 10 mg/ml in DMSO.

Description

Pentadecanoic acid is a 15:0 saturated fatty acid found in esterified form in the lipids of many bacterial species and in the milk fat of ruminants.¹ It has been used as a biological marker for the intake of dairy fat in the assessment of metabolic risk factors.^{1,2}

References

- 1. Smedman, A.E.M., Gustafsson, I.B., Berglund, L.G.T., et al. Pentadecanoic acid in serum as a marker for intake of milk fat: Relations between intake of milk fat and metabolic risk factors. Am. J. Clin. Nutr. 69(1), 22-29 (1999).
- 2. Wolk, A., Vessby, B., Ljung, H., et al. Evaluation of a biological marker of dairy fat intake. Am. J. Clin. Nutr. 68(2), 291-295 (1998).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 07/29/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM