

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Desoximetasone-d₂

Item No. 28159

Formal Name: 9-fluoro-11β,21-dihydroxy-16α-methyl-

pregna-1,4-diene-3,20-dione-d₃

Synonym: 17-deoxy Dexamethasone-d₂

MF: $C_{22}H_{26}D_3FO_4$

FW: 379.5


Chemical Purity: ≥98% (Desoximetasone)

Deuterium

Incorporation: ≥99% deuterated forms (d₁-d₃); ≤1% d₀

Supplied as: A solid Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Desoximetasone-d₃ is intended for use as an internal standard for the quantification of desoximetasone (Item No. 23945) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Desoximetasone-d₃ is supplied as a solid. A stock solution may be made by dissolving the desoximetasone-d₃ in the solvent of choice. Desoximetasone-d₃ is soluble in organic solvents such as methanol and DMSO, which should be purged with an inert gas.

Description

Desoximetasone is a glucocorticoid. In vivo, desoximetasone (0.001-0.1 mg) reverses increases in epidermal cell proliferation induced by phorbol 12-myristate 13-acetate (TPA; Item No. 10008014) in SENCAR mice when administered topically. It also reduces TPA-induced expression of IL-6, COX-2, iNOS, and c-Jun mRNA levels in the skin of SENCAR mice, indicating anti-inflammatory action. Formulations containing desoximetasone have been used in the treatment of psoriasis and atopic dermatitis.

Reference

1. Kowalczyk, P., Junco, J.J., Kowalczyk, M.C., et al. The effects of dissociated glucocorticoids RU24858 and RU24782 on TPA-induced skin tumor promotion biomarkers in SENCAR mice. Mol. Carcinog. 53(6), 488-497 (2014).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 06/06/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM