

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

# **PRODUCT** INFORMATION



### 1,2-Dilinoleoyl-3-Oleoyl-rac-glycerol

Item No. 28178

CAS Registry No.: Formal Name:	2190-21-8 (9Z,12Z)-9,12-octadecadienoic acid,	
	1,1'-[1-[[[(9Z)-1-oxo-9-octadecen-1-yl] oxy]methyl]-1,2-ethanediyl] ester	
Synonyms:	1,2-Dilinolein-3-Olein,	Q
	TG(18:2/18:2/18:1)	
MF:	C <sub>57</sub> H <sub>100</sub> O <sub>6</sub>	
FW:	881.4	
Purity:	≥95%	
Supplied as:	A liquid	 0
Storage:	-20°C	
Stability:	≥2 years	

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

#### Laboratory Procedures

1,2-Dilinoleoyl-3-oleoyl-rac-glycerol is supplied as a liquid. A stock solution may be made by dissolving the 1,2-dilinoleoyl-3-oleoyl-rac-glycerol in the solvent of choice, which should be purged with an inert gas. 1,2-dilinoleoyl-3-oleoyl-rac-glycerol is slightly soluble in methanol.

1,2-Dilinoleoyl-3-oleoyl-rac-glycerol is slightly soluble in aqueous solutions. To enhance aqueous solubility, dilute the organic solvent solution into aqueous buffers or isotonic saline. If performing biological experiments, ensure the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. We do not recommend storing the aqueous solution for more than one day.

#### Description

1,2-Dilinoleoyl-3-oleoyl-rac-glycerol is a triacylglycerol that contains linoleic acid (Item Nos. 90150 | 90150.1 | 21909) at the sn-1 and sn-2 positions and oleic acid (Item Nos. 90260 | 24659) at the sn-3 position. It has been found in ostrich and emu oils, as well as in the fat body of male B. lapidarius bumblebees.<sup>1,2</sup> 1,2-Dilinoleoyl-3-oleoyl-rac-glycerol (3% w/v) reduces scald development on apples of the Delicious variety when applied immediately following harvest and assessed after six months of storage.<sup>3</sup>

#### References

- 1. Zhou, Y., Xue, Y., Chen, G.C., et al. Rapid separation and characterisation of triacylglycerols in ostrich oil by ultra performance liquid chromatography coupled with quadrupole time-of-flight mass spectrometry. Food Chem. 141(3), 2098-2102 (2013).
- 2. Cvačka, J., Hovorka, O., Jiroš, P., et al. Analysis of triacylglycerols in fat body of bumblebees by chromatographic methods. J. Chromatogr. A. 1101(1-2), 226-237 (2006).
- 3. Ju, Z., Duan, Y., and Ju, Z. Mono-, di-, and tri-acylglycerols and phospholipids from plant oils inhibit scald development in 'Delicious' apples. Postharvest Bio. Technol. 19(1), 1-7 (2000).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 08/19/2019

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM