



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION




Bilastine

Item No. 28375

CAS Registry No.: 202189-78-4
Formal Name: 4-[2-[4-[1-(2-ethoxyethyl)-1H-benzimidazol-2-yl]-1-piperidiny]ethyl]- α,α -dimethyl-benzeneacetic acid

Synonym: F-96221-BM1
MF: $C_{28}H_{37}N_3O_3$
FW: 463.6
Purity: $\geq 98\%$
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥ 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Bilastine is supplied as a crystalline solid. A stock solution may be made by dissolving the bilastine in the solvent of choice, which should be purged with an inert gas. Bilastine is soluble in the organic solvent chloroform at a concentration of approximately 30 mg/ml.

Description

Bilastine is a histamine H_1 receptor antagonist ($IC_{50} = 180$ nM).¹ It is selective for the histamine H_1 receptor in a panel of 30 receptors *in vitro* at 100 μM . Bilastine prevents microvascular extravasation ($ED_{50} = 185$ $\mu\text{g}/\text{kg}$, i.v.), bronchospasm ($ED_{50} = 4.6$ $\mu\text{g}/\text{kg}$, i.v.), and systemic anaphylaxis ($ED_{50} = 0.2$ $\mu\text{g}/\text{kg}$, p.o.) induced by subcutaneous histamine in guinea pigs.² It prevents anaphylaxis induced by subcutaneous administration of ovalbumin or dinitrophenylated human albumin (DNP) in sensitized rats when administered at doses of 7.6 and 6.0 mg/kg, respectively. Formulations containing bilastine have been used in the treatment of urticaria and allergic rhinitis.

References

1. Corcóstegui, R., Labeaga, L., Inneráritu, A., *et al.* Preclinical pharmacology of bilastine, a new selective histamine H_1 receptor antagonist: Receptor selectivity and *in vitro* antihistaminic activity. *Drugs R.D.* **6(6)**, 371-384 (2005).
2. Corcóstegui, R., Labeaga, L., Inneráritu, A., *et al.* *In vivo* pharmacological characterisation of bilastine, a potent and selective histamine H_1 receptor antagonist. *Drugs R.D.* **7(4)**, 219-231 (2006).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/26/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM