

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Cortexolone 17a-propionate

Item No. 28392

CAS Registry No.:	19608-29-8	
Formal Name:	21-hydroxy-17-(1-oxopropoxy)-	HO
	pregn-4-ene-3,20-dione	0
Synonym:	CB-03-01	
MF:	C ₂₄ H ₃₄ O ₅	
FW:	402.5	$f \neq \backslash \rangle$
Purity:	≥98%	
UV/Vis.:	λ _{max} : 240, 295 nm	
Supplied as:	A solid	
Storage:	-20°C	0
Stability:	≥2 years	

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Cortexolone 17a-propionate is supplied as a solid. A stock solution may be made by dissolving the cortexolone 17α -propionate in the solvent of choice, which should be purged with an inert gas. Cortexolone 17α -propionate is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of cortexolone 17a-propionate in DMSO is approximately 10 mg/ml and approximately 20 mg/ml in ethanol and DMF.

Cortexolone 17α -propionate is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, cortexolone 17α -propionate should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. Cortexolone 17α -propionate has a solubility of approximately 0.1 mg/ml in a 1:9 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Cortexolone 17α -propionate is a peripherally selective and rogen receptor antagonist.¹ Cortexolone 17a-propionate binds to the androgen receptor (AR) and inhibits AR-regulated transcription in a reporter assay.² It also inhibits androgen-regulated lipid synthesis in primary human sebocytes. Cortexolone 17a-propionate inhibits inflammatory cytokine production from primary human sebocytes and scalp dermal papilla cells.^{2,3} It inhibits testosterone propionate-induced flank organ enlargement in Syrian golden hamsters by 84% when administered topically at a dose of 400 μ g/animal.

References

- 1. Celasco, G., Moro, L., Bozzella, R., et al. Biological profile of cortexolone 17α-propionate (CB-03-01), a new topical and peripherally selective androgen antagonist. Arzneimittelforschung 54(12), 881-886 (2004).
- 2. Rosette, C., Agan, F.J., Mazzetti, A., et al. Cortexolone 17α-propionate (clascoterone) is a novel androgen receptor antagonist that inhibits production of lipids and inflammatory cytokines from sebocytes in vitro. J. Drugs Dermatol. 18(5), 412-418 (2019).
- 3. Rosette, C., Rosette, N., Mazezetti, A., et al. Cortexolone 17α-propionate (clascoterone) is an androgen receptor antagonist in dermal papilla cells in vitro. J. Drugs Dermatol. 18(2), 197-201 (2019).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/29/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM