



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## (±)-Synephrine (hydrochloride)

Item No. 28398

**CAS Registry No.:** 5985-28-4  
**Formal Name:** 4-hydroxy- $\alpha$ -[(methylamino)methyl]-benzenemethanol, monohydrochloride  
**MF:** C<sub>9</sub>H<sub>13</sub>NO<sub>2</sub> • HCl  
**FW:** 203.7  
**Purity:** ≥95%  
**UV/Vis.:**  $\lambda_{\text{max}}$ : 226, 279 nm  
**Supplied as:** A crystalline solid  
**Storage:** -20°C  
**Stability:** ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

(±)-Synephrine (hydrochloride) is supplied as a crystalline solid. A stock solution may be made by dissolving the (±)-synephrine (hydrochloride) in the solvent of choice, which should be purged with an inert gas. (±)-Synephrine (hydrochloride) is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of (±)-synephrine (hydrochloride) in these solvents is approximately 30 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of (±)-synephrine (hydrochloride) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of (±)-synephrine (hydrochloride) in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

### Description

(±)-Synephrine is an alkaloid with vasoconstrictor and metabolic activities.<sup>1-3</sup> It binds to  $\alpha_{1A}$ -,  $\alpha_{2A}$ -, and  $\alpha_{2C}$ -adrenergic receptors (ARs;  $K_i$ s = 78, 36.7, and 24.4  $\mu$ M, respectively).<sup>4</sup> (±)-Synephrine is an agonist of  $\alpha_{1A}$ -ARs in HEK293 cells ( $EC_{50}$  = 4  $\mu$ M in a reporter assay) but not in CHO cells expressing  $\alpha_{2A}$ - or  $\alpha_{2C}$ -AR. It also acts as an antagonist of  $\alpha_{1A}$ -,  $\alpha_{2A}$ -, and  $\alpha_{2C}$ -ARs, inhibiting L-phenylephrine-induced activation of  $\alpha_{1A}$ -AR in HEK293 cells and activation of  $\alpha_{2A}$ - and  $\alpha_{2C}$ -ARs induced by the  $\alpha_2$ -AR agonist medetomidine in CHO cells ( $IC_{50}$ s = 12.8, 26, and 27.3  $\mu$ M, respectively, in reporter assays). (±)-Synephrine induces contractions in isolated rabbit aortic rings ( $EC_{50}$  = 6.2  $\mu$ g/ml) and increases ligation-induced mean arterial pressure in rats when administered at a dose of 2 mg/kg per day.<sup>1,5</sup> It induces lipolysis in isolated rat and human adipocytes when used at concentrations of 100 and 1,000  $\mu$ g/ml.<sup>2</sup> (±)-Synephrine (50  $\mu$ M) increases phosphorylation of Akt and AMP-activated protein kinase (AMPK) and translocation of Glut4 to the plasma membrane, as well as increases insulin-induced glucose consumption in L6 muscle cells when used at concentrations ranging from 25 to 200  $\mu$ M.<sup>3</sup>

### References

1. Huang, Y.-T., Wang, G.-F., Chen, C.-F., et al. *Life Sci.* **57(22)**, 2011-2020 (1995).
2. Mercader, J., Wanecq, E., Chen, J., et al. *J. Physiol. Biochem.* **67(3)**, 443-452 (2011).
3. Hong, N.-Y., Cui, Z.-G., Kang, H.-K., et al. *Biochem. Biophys. Res. Commun.* **418(4)**, 720-724 (2012).
4. Ma, G., Bavadekar, S.A., Schaneberg, B.T., et al. *Planta Med.* **76(10)**, 981-986 (2010).
5. Bevan, J.A. and Osher, J.V. *J. Pharmacol. Exp. Ther.* **150(3)**, 370-374 (1965).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/04/2019

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM