



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## TLR7/8 Agonist 1 (hydrochloride)

Item No. 28451

**CAS Registry No.:** 1620278-72-9  
**Formal Name:** 1-[[4-(aminomethyl)phenyl]methyl]-2-butyl-1H-imidazo[4,5-c]quinolin-4-amine, dihydrochloride  
**MF:** C<sub>22</sub>H<sub>25</sub>N<sub>5</sub> • 2HCl  
**FW:** 432.4  
**Purity:** ≥98%  
**UV/Vis.:** λ<sub>max</sub>: 229, 247, 310, 323 nm  
**Supplied as:** A solid  
**Storage:** -20°C  
**Stability:** ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

TLR7/8 agonist 1 (hydrochloride) is supplied as a solid. A stock solution may be made by dissolving the TLR7/8 agonist 1 (hydrochloride) in the solvent of choice, which should be purged with an inert gas. TLR7/8 agonist 1 (hydrochloride) is soluble in the organic solvent DMSO.

### Description

TLR7/8 agonist 1 is a dual agonist of toll-like receptor 7 (TLR7) and TLR8 (EC<sub>50</sub>s = 50 and 55 nM, respectively, in cell-based assays).<sup>1</sup> It increases the levels of TNF-α, IFN-γ, IL-12p40, and chemokine (C-C motif) ligand 4 (CCL4) in human peripheral blood mononuclear cells (PBMCs) in a biphasic manner. TLR7/8 agonist 1 (25 nmol, s.c.) increases serum levels of IL-12p40 and chemokine (C-X-C motif) ligand 10 (CXCL10), as well as the number of dendritic cells per injection-site proximal lymph node, in mice.<sup>2</sup> It has been conjugated to various fluorophores as TLR7 probes and to polymer particles for the modulation of TLR7/8 agonist 1 adjuvant activity.<sup>2,3</sup>

### References

1. Beesu, M., Caruso, G., Salyer, A.C., *et al.* Structure-based design of human TLR8-specific agonists with augmented potency and adjuvanticity. *J. Med. Chem.* **58(19)**, 7833-7849 (2015).
2. Lynn, G.M., Chytil, P., Francica, J.R., *et al.* Impact of polymer-TLR-7/8 agonist (adjuvant) morphology on the potency and mechanism of CD8 T cell induction. *Biomacromolecules* **20(2)**, 854-870 (2019).
3. Shukla, N.M., Mutz, C.A., Ukani, R., *et al.* Syntheses of fluorescent imidazoquinoline conjugates as probes of Toll-like receptor 7. *Bioorg. Med. Chem. Lett.* **20(22)**, 6384-6386 (2010).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/18/2019

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM