



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Hexanoic Acid-d₃

Item No. 28497

CAS Registry No.: 55320-69-9
Formal Name: hexanoic-6,6,6-d₃ acid
Synonyms: C6:0-d₃, Caproic Acid-d₃
MF: C₆H₉D₃O₂
FW: 119.2
Chemical Purity: ≥95% (Hexanoic Acid)
Deuterium Incorporation: ≥99% deuterated forms (d₁-d₃); ≤1% d₀
Supplied as: An oil
Storage: -20°C
Stability: ≥1 year



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Hexanoic acid-d₃ is intended for use as an internal standard for the quantification of hexanoic acid by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Hexanoic acid-d₃ is supplied as an oil. A stock solution may be made by dissolving the hexanoic acid-d₃ in the solvent of choice, which should be purged with an inert gas. Hexanoic acid-d₃ is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of hexanoic acid-d₃ in these solvents is approximately 30 mg/ml.

Description

Hexanoic acid is a short-chain saturated fatty acid that has been found in various animal fats used in biodiesel production.¹

Reference

1. Sales, E.A., Ghirardi, M.L., and Jorquera, O. Subcritical ethylic biodiesel production from wet animal fat and vegetable oils: A net energy ratio analysis. *Energy Convers. Manag.* **141**, 216-223 (2017).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/25/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM