



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION



**Ro 08-2750**  
Item No. 28772

**CAS Registry No.:** 37854-59-4  
**Formal Name:** 2,3,4,10-tetrahydro-7,10-dimethyl-2,4-dioxo-benzo[g]pteridine-8-carboxaldehyde  
**MF:** C<sub>13</sub>H<sub>10</sub>N<sub>4</sub>O<sub>3</sub>  
**FW:** 270.2  
**Purity:** ≥95%  
**UV/Vis.:** λ<sub>max</sub>: 248, 288, 335, 460 nm  
**Supplied as:** A solid  
**Storage:** -20°C  
**Stability:** ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

## Laboratory Procedures

Ro 08-2750 is supplied as a solid. A stock solution may be made by dissolving the Ro 08-2750 in the solvent of choice, which should be purged with an inert gas. Ro 08-2750 is soluble in the organic solvent DMSO.

## Description

Ro 08-2750 is a p75 neurotrophin receptor (p75<sup>NTR</sup>) ligand that completely inhibits nerve growth factor (NGF) binding to p75<sup>NTR</sup> in PC12 cells when used at a concentration of 100 μM.<sup>1</sup> It also binds to NGF dimers (K<sub>d</sub> = ~1 μM) to induce a conformational change that prevents NGF binding to p75<sup>NTR</sup> or tropomyosin-related kinase A (TrkA). Ro 08-2750 inhibits NGF-induced death of SK-N-MC cells and inhibits autophosphorylation of TrkA in PC12 cells. It reverses NGF-induced decreases in β-catenin mRNA expression in, and migration of, SKOV3 and OVCAR-3 ovarian cancer cells.<sup>2</sup>

## References

1. Niederhauser, O., Mangold, M., Schubeneil, R., *et al.* NGF ligand alters NGF signaling via p75<sup>NTR</sup> and trkA. *J. Neurosci. Res.* **61(3)**, 263-272 (2000).
2. Li, B., Cai, S., Zhao, Y., *et al.* Nerve growth factor modulates the tumor cells migration in ovarian cancer through the WNT/β-catenin pathway. *Oncotarget* **7(49)**, 81026-81048 (2016).

### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/02/2019

## CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM