



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



PAD4 (mouse, recombinant)


Item No. 28910

Overview and Properties

Synonyms:	PADI4, PADI5, Peptidylarginine Deiminase 4, Protein Arginine Deiminase Type 4, Protein Arginine Deiminase Type IV
Source:	Active recombinant N-terminal His-tagged protein expressed in insect cells
Amino Acids:	2-666 AA (full length)
Uniprot No.:	Q9Z183
Molecular Weight:	76.08 kDa
Storage:	-80°C (as supplied); avoid freeze/thaw cycles by aliquoting protein
Stability:	≥1 year
Purity:	≥90% estimated by SDS-PAGE
Supplied in:	50 mM Tris HCl, pH 8.0, with 500 mM NaCl and 5% glycerol
Protein Concentration:	<i>batch specific</i> mg/ml
Activity:	<i>batch specific</i> U/ml
Specific Activity:	<i>batch specific</i> U/mg
Unit Definition:	One unit is defined as the amount of enzyme to produce 1 nmol of NH ₄ ⁺ per minute at 37°C in 50 mM HEPES, pH 7.7, containing 10 mM calcium chloride, 5 mM DTT, and 2 mM N-α-benzoyl-L-arginine ethyl ester (BAEE).


Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Images



Lane 1: MW Markers
Lane 2: PAD4 (2 µg)
Lane 3: PAD4 (4 µg)

Representative gel image shown; actual purity may vary between each batch.



WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/01/2021

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Description

Protein arginine deiminase 4 (PAD4) catalyzes the conversion of arginine residues to citrulline within cellular protein substrates, resulting in the loss of a positive charge, which can alter protein structure and/or function.¹ It is expressed in neutrophils, as well as a variety of tissues, including the brain, liver, lung, and kidney.^{1,2,5} PAD4 has a key role in NETosis, a lytic form of cell death characterized by the release of neutrophil extracellular traps (NETs).¹ Upon neutrophil activation, PAD4 translocates to the nucleus where it citrullinates histones, initiating chromatin decondensation and the release of NETs.^{2,3,6} Neutrophils isolated from *Pad4*^{-/-} mice exhibit decreased citrullination of histone H3 under both basal and LPS-stimulated conditions and are defective for NET formation in response to stimulation with LPS, phorbol 12-myristate 13-acetate (PMA; Item No. 10008014), or hydrogen peroxide.³ *Pad4*^{-/-} mice exhibit larger lesions than wild-type mice in a model of necrotizing fasciitis induced by M1 group A *S. pyogenes* lacking the extracellular DNase Sda1. *Pad4*-deficient mice also exhibit reduced infarct size in a model of myocardial ischemia-reperfusion injury and reduced tumor growth in a Lewis lung carcinoma model.^{2,4} Serum PAD4 autoantibodies have been found in patients with rheumatoid arthritis and are associated with disease severity. Cayman's PAD4 (mouse, recombinant) can be used for ELISA, enzyme activity assay, and Western blot (WB) applications.

References

1. van Beers, J.J.B.C., Zendman, A.J.W., Raijmakers, R., *et al.* Peptidylarginine deiminase expression and activity in PAD2 knock-out and PAD4-low mice. *Biochimie* **95(2)**, 299-308 (2013).
2. Demers, M., Wong, S.L., Martinod, K., *et al.* Priming of neutrophils toward NETosis promotes tumor growth. *Oncoimmunology* **5(5)**, e1134073 (2016).
3. Li, P., Li, M., Lindberg, M.R., *et al.* PAD4 is essential for antibacterial innate immunity mediated by neutrophil extracellular traps. *J. Exp. Med.* **207(9)**, 1853-1862 (2010).
4. Savchenko, A.S., Borissoff, J.I., Martinod, K., *et al.* VWF-mediated leukocyte recruitment with chromatin decondensation by PAD4 increases myocardial ischemia/reperfusion injury in mice. *Blood* **123(1)**, 141-148 (2014).
5. Jones, J.E., Causey, C.P., Knuckley, B., *et al.* Protein arginine deiminase 4 (PAD4): Current understanding and future therapeutic potential. *Curr. Opin. Drug Discov. Devel.* **12(5)**, 616-627 (2009).
6. Thiam, H.R., Wong, S.L., Qiu, R., *et al.* NETosis proceeds by cytoskeleton and endomembrane disassembly and PAD4-mediated chromatin decondensation and nuclear envelope rupture. *Proc. Natl. Acad. Sci. USA* **117(13)**, 7326-7337 (2020).

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM