

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



N-Oleoyl Glutamine

Item No. 28922

CAS Registry No.: 247150-73-8

Formal Name: N^2 -[(9Z)-1-oxo-9-octadecen-1-yl]-

L-glutamine

MF: $C_{23}H_{42}N_2O_4$ FW: 410.6 **Purity:** ≥95%

Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

N-Oleoyl glutamine is supplied as a crystalline solid. A stock solution may be made by dissolving the N-Oleoyl glutamine in the solvent of choice, which should be purged with an inert gas. N-Oleoyl glutamine is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of N-oleoyl glutamine in ethanol is approximately 1 mg/ml and approximately 30 mg/ml in DMSO and DMF.

Description

N-Oleoyl glutamine is an endogenous N-acyl amine. 1.2 It is both formed and degraded by peptidase M20 domain-containing 1 (PM20D1), a factor secreted by thermogenic adipose cells. N-Oleoyl glutamine (50 µM) induces mitochondrial uncoupling to increase respiration in C2C12 cells by 64%.³ It inhibits capsaicin-induced calcium flux in HEK293A cells co-transfected with transient receptor potential vanilloid 1 (TRPV1) and red fluorescent protein (RFP).² N-Oleoyl glutamine (100 mg/kg) reduces the number of acetic acid-induced abdominal constrictions, as well as formalin-induced paw licking in mice.

References

- 1. Long, J.Z., Svensson, K.J., Bateman, L.A., et al. The secreted enzyme PM20D1 regulates lipidated amino acid uncouplers of mitochondria. Cell 166(2), 1-12 (2016).
- 2. Long, J.Z., Roche, A.M., Berdan, C.A., et al. Ablation of PM20D1 reveals N-acyl amino acid control of metabolism and nociception. Proc. Natl. Acad. Sci. U.S.A. 115(29), E6937-E6945 (2018).
- 3. Lin, H., Long, J.Z., Roche, A.M., et al. Discovery of hydrolysis-resistant isoindoline N-acyl amino acid analogues that stimulate mitochondrial respiration. J. Med. Chem. 61(7), 3224-3230 (2018)

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 07/24/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM