

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



1,2-Dioleoyl-3-Arachidoyl-rac-glycerol

Item No. 29060

| CAS Registry No.: Formal Name: | 77145-65-4 eicosanoic acid, 2,3- <i>bis</i> [[(9Z)-1-oxo-9- | 0 |
|--|---|---|
| Synonyms: | octadecen-1-yl]oxy]propyl ester 1,2-Olein-3-Arachidin, TG(18:1/18:1/20:0) | |
| MF: | C ₅₉ H ₁₁₀ O ₆ | |
| FW: | 915.5 | |
| Purity: | ≥95% | |
| Supplied as: | A crystalline solid | $\qquad \qquad $ |
| Storage: | -20°C | 0 |
| Stability: | ≥2 years | |
| Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis. | | |

Laboratory Procedures

1,2-Dioleoyl-3-arachidoyl-rac-glycerol is supplied as a crystalline solid. A stock solution may be made by dissolving the 1,2-dioleoyl-3-arachidoyl-rac-glycerol in the solvent of choice, which should be purged with an inert gas. 1,2-Dioleoyl-3-arachidoyl-rac-glycerol is soluble in organic solvents such as ethanol and dimethyl formamide. The solubility of 1,2-dioleoyl-3-arachidoyl-rac-glycerol in these solvents is approximately 10 mg/ml. It is also slightly soluble in chloroform.

1,2-Dioleoyl-3-arachidoyl-rac-glycerol is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, 1,2-dioleoyl-3-arachidoyl-rac-glycerol should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. 1,2-Dioleoyl-3-arachidoyl-rac-glycerol has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

1,2-Dioleoyl-3-arachidoyl-*rac*-glycerol is a triacylglycerol that contains oleic acid (Item Nos. 90260 | 24659) at the sn-1 and sn-2 positions and arachidic acid (Item Nos. 9000339 | 21906) at the sn-3 position. It has been found in hazelnut, cashew, date seed, corn, sunflower, and soybean oils.^{1,2}

References

- 1. Holcapek, M., Lísa, M., Jandera, P., et al. Quantitation of triacylglycerols in plant oils using HPLC with APCI-MS, evaporative light-scattering, and UV detection. J. Sep. Sci. 28(12), 1315-1333 (2005).
- 2. Gao, B., Luo, Y., Lu, W., et al. Triacylglycerol compositions of sunflower, corn and soybean oils examined with supercritical CO₂ ultra-performance convergence chromatography combined with quadrupole time-of-flight mass spectrometry. Food Chem. 218, 569-574 (2017).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/05/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM