

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



# Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



# (-)-Nicotine

Item No. 29138

CAS Registry No.: 54-11-5

Formal Name: 3-[(2S)-1-methyl-2-pyrrolidinyl]-pyridine

Synonyms: L-Nicotine, (-)-(S)-Nicotine

MF:  $C_{10}H_{14}N_2$ FW: 162.2 **Purity:** ≥95% UV/Vis.:  $\lambda_{max}$ : 262 nm

Supplied as: An oil Storage: -20°C Stability: ≥2 years Item Origin: Synthetic





(-)-Nicotine is supplied as an oil. A stock solution may be made by dissolving the (-)-nicotine in the solvent of choice, which should be purged with an inert gas. (-)-Nicotine is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of (-)-nicotine in ethanol and DMF is approximately 50 mg/ml and approximately 30 mg/ml in DMSO.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of (-)-nicotine can be prepared by directly dissolving the oil in aqueous buffers. The solubility of (-)-nicotine in PBS, pH 7.2, is approximately 1 mg/ml. We do not recommend storing the aqueous solution for more than one day.

### Description

(-)-Nicotine is an alkaloid that has been found in tobacco. 1 It is an agonist at neuronal nicotinic acetylcholine receptors (nAChRs) and binds to α3β4 and α4β2 subunit-containing nAChRs with K, values of 481 and 11.1 nM, respectively.<sup>1,2</sup> Chronic exposure to (-)-nicotine results in increased expression of certain nAChRs, particularly α4β2 subunit-containing nAChRs.<sup>3</sup> (-)-Nicotine has addictive properties.<sup>2,3</sup> Formulations containing (-)-nicotine have been used as smoking cessation aids for the relief of nicotine withdrawal symptoms.

### References

- 1. Clayton, P., Lu, A., and Bishop, L. The pyrolysis of (-)-(S)-nicotine: Racemization and decomposition. Chirality 22(4), 442-446 (2010).
- 2. Zaveri, N., Jiang, F., Olsen, C., et al. Novel α3β4 nicotinic acetylcholine receptor-selective ligands. Discovery, structure-activity studies, and pharmacological evaluation. J. Med. Chem. 53(22), 8187-8191
- 3. Albuquerque, E.X., Pereira, E.F.R., Alkondon, M., et al. Mammalian nicotinic acetylcholine receptors: From structure to function. Physiol. Rev. 89(1), 73-120 (2009).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 11/06/2019

## **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

**FAX:** [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM