

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



3-O-methyl Quercetin

Item No. 29333

CAS Registry No.: 1486-70-0

Formal Name: 2-(3,4-dihydroxyphenyl)-5,7-dihydroxy-3-

methoxy-4H-1-benzopyran-4-one

Synonyms: NSC 154016, Quercetin 3-methyl ether

MF: $C_{16}H_{12}O_7$ 316.3 FW: **Purity:** ≥95% Supplied as: A solid Storage: -20°C Stability: ≥2 years Item Origin: Synthetic

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

3-O-methyl Quercetin is supplied as a solid. A stock solution may be made by dissolving the 3-O-methyl quercetin in the solvent of choice, which should be purged with an inert gas. 3-O-methyl Quercetin is slightly soluble in DMSO.

Description

3-O-methyl Quercetin is a flavone that has been found in C. balchaschensis and has diverse biological activities. $^{1\text{--}4}$ It is an inhibitor of β -secretase (IC $_{50}$ = 6.5 μM). 1 It scavenges DPPH (Item No. 14805) radicals and superoxide anions in cell-free assays (IC $_{50}$ s = 14.17 and 17.39 μM , respectively), as well as inhibits lipid peroxidation and hydrogen peroxide-induced neuronal cell death in primary rat cortical cells (IC₅₀s = 19 and 3.5 μM, respectively).² 3-O-methyl Quercetin inhibits poliovirus RNA synthesis in infected HeLa cells when used at concentrations of 2 and 20 µg/ml.³ It also inhibits LPS-induced nitric oxide (NO) production in BV-2 microglia (IC_{50} = 3.8 μ M) and reduces LPS-induced increases in the levels of inducible nitric oxide synthase (iNOS) in BV-2 microglia.4

References

- 1. Zhumanova, K., Lee, G., Baiseitova, A., et al. Inhibitory mechanism of O-methylated quercetins, highly potent β-secretase inhibitors isolated from Caragana balchaschensis (Kom.) Pojark. J. Ethnopharmacol. 272, 113936 (2021).
- 2. Kim, S.H., Kumar, C.N., Kim, H.J., et al. Glucose-containing flavones—their synthesis and antioxidant and neuroprotective activities. Bioorg. Med. Chem. Lett. 19(21), 6009-6013 (2009).
- Castrillo, J.L. and Carrasco, L. Action of 3-methylquercetin on poliovirus RNA replication. J. Virol. 61(10), 3319-3321 (1987).
- Kim, J.Y., Lim, H.J., and Ryu, J.-H. In vitro anti-inflammatory activity of 3-O-methyl-flavones isolated from Siegesbeckia glabrescens. Bioorg. Med. Chem. Lett. 18(4), 1511-1514 (2008).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the m can be found on our website.

Copyright Cayman Chemical Company, 08/26/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM