



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION



BMS 204352

Item No. 29378

CAS Registry No.: 187523-35-9

Formal Name: (3S)-3-(5-chloro-2-methoxyphenyl)-
(3S)-3-(5-chloro-2-methoxyphenyl)-
3-fluoro-1,3-dihydro-6-

(trifluoromethyl)-2H-indol-2-one

Synonym: (+)-BMS 204352

MF: C₁₆H₁₀ClF₄NO₂

FW: 359.7


Purity: ≥98%

UV/Vis.: λ_{max}: 211 nm

Supplied as: A crystalline solid

Storage: -20°C

Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

BMS 204352 is supplied as a crystalline solid. A stock solution may be made by dissolving the BMS 204352 in the solvent of choice, which should be purged with an inert gas. BMS 204352 is soluble in the organic solvent DMSO.

Description

BMS 204352 is an activator of large-conductance calcium-activated potassium (K_{Ca}1.1/BK) channels that increases K_{Ca}1.1 currents in *X. laevis* oocytes expressing human K_{Ca}1.1 in a calcium- and concentration-dependent manner.¹ It decreases electrically induced population excitatory postsynaptic potentials (pEPSPs) *in vitro* in CA1 rat neurons and in anesthetized rats when administered at doses ranging from 0.005 to 1 mg/kg. BMS 204352 (0.001 and 0.3 mg/kg) reduces infarct volume in a rat model of ischemic stroke induced by middle cerebral artery occlusion (MCAO). It decreases edema in the ipsilateral hippocampus, thalamus, and adjacent cortex and time to find the platform in the Morris water maze in a rat model of fluid percussion-induced traumatic brain injury (TBI).² BMS 204352 also reverses cortical hyperexcitability and reduces hyperactivity and grooming behaviors in the *Fmr1*^{-/-} mouse model of Fragile X syndrome.³ Formulations containing BMS 204352 have been used in the treatment of ischemic stroke.

References

1. Gribkoff, V.K., Starrett, J.E., Jr., Dworetzky, S.I., et al. Targeting acute ischemic stroke with a calcium-sensitive opener of maxi-K potassium channels. *Nat. Med.* **7**(4), 471-477 (2001).
2. Cheney, J.A., Weisser, J.D., Bareyre, F.M., et al. The maxi-K channel opener BMS-204352 attenuates regional cerebral edema and neurologic motor impairment after experimental brain injury. *J. Cereb. Blood Flow Metab.* **21**(4), 396-403 (2001).
3. Carreno-Munoz, M.I., Martins, F., Medrano, M.C., et al. Potential involvement of impaired BK_{Ca} channel function in sensory defensiveness and some behavioral disturbances induced by unfamiliar environment in a mouse model of fragile X syndrome. *Neuropsychopharmacology* **43**(3), 492-502 (2018).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 - USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM