

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



# Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

# SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



## Octanoic Acid-13C

Item No. 29461

CAS Registry No.: 59669-16-8 Formal Name: octanoic-1-13C acid Caprylic Acid-13C Synonym: MF:  $C_7[^{13}C]H_{16}O_2$ 

FW: 145.2 **Purity:** ≥98% Supplied as: A liquid Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

#### **Laboratory Procedures**

Octanoic acid-13C is supplied as a liquid. A stock solution may be made by dissolving the octanoic acid-13C in the solvent of choice, which should be purged with an inert gas. Octanoic acid-13C is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of octanoic acid-13C in these solvents is approximately 30 mg/ml.

## Description

Octanoic acid-13C is intended for use as an internal standard for the quantification of octanoic acid by GC- or LC-MS. Octanoic acid is a medium-chain saturated fatty acid. It has been found in Teleme cheeses made from goat, ovine, or bovine milk.<sup>1</sup> Octanoic acid is active against the bacteria S. mutans, S. gordonii, F. nucleatum, and P. gingivalis (IC $_{80}$ s = <125, <125, 1,403, and 2,294  $\mu$ M, respectively). Levels of octanoic acid are increased in the plasma of patients with medium-chain acyl-CoA dehydrogenase (MCAD) deficiency, an inborn error of fatty acid metabolism characterized by hypoketotic hypoglycemia, medium-chain dicarboxylic aciduria, and intolerance to fasting.<sup>3,4</sup>

## References

- 1. Mallatou, H., Pappa, E., and Massouras, T. Changes in free fatty acids during ripening of Teleme cheese made with ewes', goats', cows' or a mixture of ewes' and goats' milk. Int. Dairy J. 13(1-3), 211-219 (2003).
- 2. Hyang, C.B., Alimova, Y., Myers, T.M., et al. Short- and medium-chain fatty acids exhibit antimicrobial activity for oral microorganisms. Arch. Oral Biol. 56(7), 650-654 (2011).
- Onkenhout, W., Venizelos, V., van der Poel, P.F.H., et al. Identification and quantification of intermediates of unsaturated fatty acid metabolism in plasma of patients with fatty acid oxidation disorders. Clin. Chem. 41(10), 1467-1474 (1995).
- 4. Rinaldo, P., O'Shea, J.J., Coates, P.M., et al. Medium-chain acyl-CoA dehydrogenase deficiency. Diagnosis by stable-isotope dilution measurement of urinary n-hexanoylglycine and 3-phenylpropionylglycine. N. Engl. J. Med. 319(20), 1308-1313 (1988).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/02/2020

## **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM