

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Tasosartan

Item No. 29495

CAS Registry No.: 145733-36-4

Formal Name: 5,8-dihydro-2,4-dimethyl-8-[[2'-

(2H-tetrazol-5-yl)[1,1'-biphenyl]-

4-yl]methyl]-pyrido[2,3-d]

pyrimidin-7(6H)-one

Synonym: WAY-ANA-756 MF: $C_{23}H_{21}N_7O$ 411.5 FW:

Purity: ≥98% UV/Vis.:

 λ_{max} : 251 nm A crystalline solid Supplied as:

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Tasosartan is supplied as a crystalline solid. A stock solution may be made by dissolving the tasosartan in the solvent of choice, which should be purged with an inert gas. Tasosartan is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of tasosartan in these solvents is approximately 2, 14, and 20 mg/ml, respectively.

Tasosartan is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, tasosartan should first be dissolved in DMF and then diluted with the aqueous buffer of choice. Tasosartan has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Tasosartan is an angiotensin II type 2 (AT₂) receptor antagonist (IC₅₀ = 1.2 nM in rat adrenal gland membranes). It also binds to the AT_1 receptor ($K_1 = 46.6$ nM in rat liver epithelial cells). Tasosartan (3 mg/kg, intraduodenally) reduces the AT₂-induced pressor response in rats.³ It also decreases mean arterial pressure in a rat model of hypertension induced by renal artery constriction when administered intragastrically or intravenously at doses of 1 and 3 mg/kg.¹

References

- 1. Ellingboe, J.W., Collini, M.D., Quagliato, D., et al. Metabolites of the angiotensin II antagonist tasosartan: The importance of a second acidic group. J. Med. Chem. 41(22), 4251-4260 (1998).
- 2. Hines, J., Fluharty, S.J., and Sakai, R.R. The angiotensin AT₁ receptor antagonist irbesartan has near-peptide affinity and potently blocks receptor signaling. Eur. J. Pharmacol. 384(1), 81-89 (1999).
- 3. Ellingboe, J.W., Antane, M., Nguyen, T.T., et al. Pyrido[2,3-d]pyrimidine angiotensin II antagonists. J. Med. Chem. 37(4), 542-550 (1994).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 02/17/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM