



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## (±)-4-CMTB

Item No. 29680

**CAS Registry No.:** 300851-67-6  
**Formal Name:** 4-chloro- $\alpha$ -(1-methylethyl)-N-2-thiazolyl-benzeneacetamide  
**MF:** C<sub>14</sub>H<sub>15</sub>ClN<sub>2</sub>OS  
**FW:** 294.8  
**Purity:**  $\geq$ 95%  
**UV/Vis.:**  $\lambda_{\text{max}}$ : 223, 269 nm  
**Supplied as:** A solid  
**Storage:** -20°C  
**Stability:**  $\geq$ 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

(±)-4-CMTB is supplied as a solid. A stock solution may be made by dissolving the (±)-4-CMTB in the solvent of choice, which should be purged with an inert gas. (±)-4-CMTB is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of (±)-4-CMTB in these solvents is approximately 3, 5, and 0.3 mg/ml, respectively.

(±)-4-CMTB is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, (±)-4-CMTB should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. (±)-4-CMTB has a solubility of approximately 0.3 mg/ml in a 1:2 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

### Description

(±)-4-CMTB is a positive allosteric modulator of free fatty acid receptor 2 (FFAR2/GPR43) with an EC<sub>50</sub> value of 0.42  $\mu$ M for [<sup>35</sup>S]GTP $\gamma$ S binding to 293 T-REx cell membranes expressing the human receptor.<sup>1</sup> It selectively induces ERK1/2 phosphorylation in 293 T-REx cells expressing human FFAR2, but not FFAR1 or FFAR3.

### Reference

1. Smith, N.J., Ward, R.J., Stoddart, L.A., *et al.* Extracellular loop 2 of the free fatty acid receptor 2 mediates allosterism of a phenylacetamide ago-allosteric modulator. *Mol. Pharmacol.* **80**(1), 163-173 (2011).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 04/10/2020

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM