



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION




Epimedin B

Item No. 29694

CAS Registry No.: 110623-73-9
Formal Name: 3-[[6-deoxy-2-O-β-D-xylopyranosyl-α-L-mannopyranosyl]oxy]-7-(β-D-glucopyranosyloxy)-5-hydroxy-2-(4-methoxyphenyl)-8-(3-methyl-2-buten-1-yl)-4H-1-benzopyran-4-one

MF: C₃₈H₄₈O₁₉
FW: 808.8
Purity: ≥98%
UV/Vis.: λ_{max}: 272, 316 nm
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Epimedin B is supplied as a crystalline solid. A stock solution may be made by dissolving the epimedin B in the solvent of choice, which should be purged with an inert gas. Epimedin B is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of epimedin B in these solvents is approximately 1, 30, and 15 mg/ml, respectively.

Epimedin B is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, epimedin B should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. Epimedin B has a solubility of approximately 0.14 mg/ml in a 1:6 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Epimedin B is a flavonoid glycoside that has been found in *E. koreanum* and has antioxidant, anti-osteoporotic, and anti-inflammatory activities.¹⁻⁵ It scavenges 2,2-diphenyl-1-picrylhydrazyl (DPPH; Item No. 14805) radicals in a cell-free assay (IC₅₀ = 107.9 μM).² Epimedin B increases proliferation of UMR-106 osteoblasts by 24.9 and 36.3% when used at concentrations of 10 and 100 μM, respectively.³ It prevents osteoporosis induced by prednisolone (Item No. 20866) in zebrafish when used at a concentration of 1 μM.⁴ Epimedin B inhibits ear edema induced by phorbol 12-myristate 13-acetate (TPA; Item No. 10008014) in mice (ED₅₀ = 114 nmol/ear).⁵

References

1. Ito, Y., Hirayama, F., Suto, K., *et al.* *Phytochemistry* **27**(3), 911-913 (1988).
2. Kim, E.S., Kim, M.K., Kang, H.K., *et al.* *Nat. Prod. Sci.* **14**(4), 233-238 (2008).
3. Meng, F.-H., Li, Y.-B., Xiong, A.-L., *et al.* *Phytomedicine* **12**(3), 189-93 (2005).
4. Zhan, Y., Wei, Y.-J., Wang, C.-M., *et al.* *J. Chin. Pharma. Sci.* **49**(1), 30-35 (2014).
5. Yasukawa, K., Ko, S.-K., and Whang, W.-K. *J. Pharm. Nutr. Sci.* **6**(2), 38-42 (2016).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/24/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM