



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Eact

Item No. 29791

CAS Registry No.: 461000-66-8
Formal Name: 3,4,5-trimethoxy-N-(2-methoxyethyl)-N-(4-phenyl-2-thiazolyl)-benzamide
MF: C₂₂H₂₄N₂O₅S
FW: 428.5
Purity: ≥98%
UV/Vis.: λ_{max}: 272 nm
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Eact is supplied as a crystalline solid. A stock solution may be made by dissolving the Eact in the solvent of choice, which should be purged with an inert gas. Eact is soluble in the organic solvent DMSO.

Description

Eact is an activator of the calcium-activated chloride channel anoctamin 1 (ANO1).¹ It increases chloride currents in FRT cells expressing human ANO1 (EC₅₀ = 3 μM). It is selective for ANO1 over cystic fibrosis transmembrane conductance regulator (CFTR) chloride channels and epithelial sodium channels (ENaCs) in FRT cells at 10 μM. Eact (20 μM) increases the rate of submucosal gland fluid secretion in human bronchi isolated from patients with cystic fibrosis and healthy individuals. It increases production of mitochondrial reactive oxygen species (ROS), reduces the mitochondrial membrane potential, and induces apoptosis in rat lung microvascular endothelial cells (RLMVECs), as well as reduces total cell counts in pulmonary artery endothelial cell populations isolated from patients with idiopathic pulmonary arterial hypertension (IPAH).² Eact also activates transient receptor potential vanilloid 1 (TRPV1) in HEK293T cells expressing the mouse receptor (EC₅₀ = 11.6 μM) and induces scratching and licking behaviors in wild-type, but not *Trpv1*^{-/-}, mice.³

References

1. Namkung, W., Yao, Z., Finkbeiner, W.E., *et al.* Small-molecule activators of TMEM16A, a calcium-activated chloride channel, stimulate epithelial chloride secretion and intestinal contraction. *FASEB J.* **25**(11), 4048-4062 (2011).
2. Allawzi, A.M., Vang, A., Clements, R.T., *et al.* Activation of anoctamin-1 limits pulmonary endothelial cell proliferation via p38-mitogen-activated protein kinase-dependent apoptosis. *Am. J. Respir. Cell Mol. Biol.* **58**(5), 658-667 (2018).
3. Liu, S., Feng, J., Luo, J., *et al.* Eact, a small molecule activator of TMEM16A, activates TRPV1 and elicits pain- and itch-related behaviours. *Br. J. Pharmacol.* **173**(7), 1208-1218 (2016).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/26/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM