

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Otilonium (bromide)

Item No. 29907

CAS Registry No.:	26095-59-0	
Formal Name:	N,N-diethyl-N-methyl-2-[[4-[[2-	
	(octyloxy)benzoyl]amino]benzoyl] oxy]-ethanaminium, monobromide	0 J
Synonym:	Octylonium	
MF:	$C_{29}H_{43}N_2O_4 \bullet Br$	
FW:	563.6	
Purity:	≥98%	•Br
UV/Vis.:	λ _{max} : 292 nm	
Supplied as:	A crystalline solid	
Storage:	-20°C	
Stability:	≥2 years	

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Otilonium (bromide) is supplied as a crystalline solid. A stock solution may be made by dissolving the otilonium (bromide) in the solvent of choice, which should be purged with an inert gas. Otilonium (bromide) is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of otilonium (bromide) in these solvents is approximately 10, 25, and 20 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of otilonium (bromide) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of otilonium (bromide) in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Otilonium is an inhibitor of L-type (IC₅₀ = 2.3 μ M) and T-type calcium channels (IC₅₀s = 0.8, 1.1, and 0.4 μ M for Ca_v3.1, Ca_v3.2, and Ca_v3.3, respectively).^{1,2} It is also an antagonist of muscarinic acetylcholine receptors (mAChRs; IC₅₀s = 0.054, 0.4, 0.222, and 0.156 μ M for M₁, M₂, M₄, and M₅ muscarinic receptors, respectively) and the platelet activating factor (PAF) receptor (IC₅₀ = 0.0552 μ M).³ Otilonium inhibits the inward calcium current in isolated rat colonic smooth muscle cells (EC₅₀ = 885 nM), an effect that can be blocked by the L-type calcium channel inhibitor nifedipine (Item No. 11106).¹ It inhibits contractions induced by the muscarinic acetylcholine receptor (mAChR) agonist methacholine in isolated circular muscle of the guinea pig proximal colon (IC₅₀ = 3.7 μ M).⁴ Otilonium (4 mg/kg) decreases fecal excretion and wet dog shakes and increases the tail withdrawal latency in the tail-flick test in a rat model of opioid withdrawal syndrome induced by morphine and naloxone.⁵

References

- 1. Martin, M.T., Hove-Madsen, L., and Jimenez, M. Neurogastroenterol. Motil. 16(2), 175-183 (2004).
- 2. Strege, P.R., Sha, L., Beyder, A., et al. Am. J. Physiol. Gastrointest. Liver Physiol. 298(5), G706-G713 (2010).
- 3. Evangelista, S., Giachetti, A., Chapelain, B., et al. Pharmacol. Res. 38(2), 111-117 (1998).
- Santicioli, P., Zagorodnyuk, V., Renzetti, A.R., et al. Naunyn-Schmiedebergs Arch. Pharmacol. 359(5), 4. 420-427 (1999).
- 5. Pinelli, A., Trivulzio, S., and Vignati, S. Toxicology 122(1-2), 23-37 (1997).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 03/27/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM