

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



ML-SA1

Item No. 29958

CAS Registry No.: 332382-54-4

Formal Name: 2-[2-(3,4-dihydro-2,2,4-trimethyl-

1(2H)-quinolinyl)-2-oxoethyl]-1H-

isoindole-1,3(2H)-dione

Synonym: Mucolipin Synthetic Agonist 1

MF: $C_{22}H_{22}N_2O_3$ FW: 362.4 **Purity:** ≥98%

UV/Vis.: λ_{max} : 216 nm Supplied as: A crystalline solid

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

ML-SA1 is supplied as a crystalline solid. A stock solution may be made by dissolving the ML-SA1 in the solvent of choice, which should be purged with an inert gas. ML-SA1 is soluble in the organic solvent dimethyl formamide (DMF) at a concentration of approximately 25 mg/ml.

ML-SA1 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, ML-SA1 should first be dissolved in DMF and then diluted with the aqueous buffer of choice. ML-SA1 has a solubility of approximately 0.2 mg/ml in a 1:4 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

ML-SA1 is an agonist of transient receptor potential mucolipin 1 (TRPML1).¹ It induces cytosolic calcium increases in HEK293T cells expressing surface-expressed mutant TRPML1 channels when used at a concentration of 10 μ M. ML-SA1 activates TRPML1-mediated current (I_{ML1}) in isolated vacuoles that were enlarged by vacuolin-1 (Item No. 20425) prior to isolation from HEK293 cells expressing TRPML1. It also activates whole-endolysosome I_{ML1} in primary murine macrophages.

Reference

1. Shen, D., Wang, X., Li, X., et al. Lipid storage disorders block lysosomal trafficking by inhibiting a TRP channel and lysosomal calcium release. Nat. Commun. 3, 731 (2012).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 05/19/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM