



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Domiphen (bromide)

Item No. 30276

CAS Registry No.: 538-71-6
Formal Name: N,N-dimethyl-N-(2-phenoxyethyl)-1-dodecanaminium, monobromide
Synonym: NSC 39415
MF: C₂₂H₄₀NO • Br
FW: 414.5
Purity: ≥98%
UV/Vis.: λ_{max}: 269 nm
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Domiphen (bromide) is supplied as a crystalline solid. A stock solution may be made by dissolving the domiphen (bromide) in the solvent of choice, which should be purged with an inert gas. Domiphen (bromide) is soluble in organic solvents such as DMSO and methanol. It is also soluble in water. We do not recommend storing the aqueous solution for more than one day.

Description

Domiphen is a quaternary ammonium compound and cationic surfactant with antimicrobial activity.^{1,2} It is active against *A. viscosus*, *A. naeslundii*, *S. mutans*, *E. coli*, and *L. monocytogenes* bacteria and inhibits *C. neoformans* yeast growth and spore germination.¹⁻⁴ It also inhibits the human-ether-a-go-go-related gene (hERG) channel (IC₅₀ = 1.5 μM in a whole-cell patch-clamp assay).⁵ Formulations containing domiphen have been used as antiseptics, disinfectants, and biocides in industrial, agricultural, veterinary, and clinical applications.

References

1. Baker, P.J., Coburn, R.A., Genco, R.J., *et al.* The in vitro inhibition of microbial growth and plaque formation by surfactant drugs. *J. Periodontal Res.* **13(5)**, 474-485 (1978).
2. Müller, A., Rychli, K., Zaiser, A., *et al.* The *Listeria monocytogenes* transposon *Tn6188* provides increased tolerance to various quaternary ammonium compounds and ethidium bromide. *FEMS Microbiol. Lett.* **361(2)**, 166-173 (2014).
3. Holdsworth, S.R. and Law, C.J. The major facilitator superfamily transporter MdtM contributes to the intrinsic resistance of *Escherichia coli* to quaternary ammonium compounds. *J. Antimicrob. Chemother.* **68(4)**, 831-839 (2013).
4. Ortiz, S.C., Huang, M., and Hull, C.M. Spore germination as a target for antifungal therapeutics. *Antimicrob. Agents Chemother.* **63(12)**, e00994-19 (2019).
5. Xia, M., Shahane, S.A., Huang, R., *et al.* Identification of quaternary ammonium compounds as potent inhibitors of hERG potassium channels. *Toxicol. Appl. Pharmacol.* **252(3)**, 250-258 (2011)

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/18/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM