



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## Clorprenaline (hydrochloride)

Item No. 30360

**CAS Registry No.:** 6933-90-0  
**Formal Name:** 2-chloro- $\alpha$ -[[[(1-methylethyl)amino]methyl]-benzenemethanol, monohydrochloride  
**Synonyms:** Isoprophenamine, NSC 334693  
**MF:** C<sub>11</sub>H<sub>16</sub>ClNO • HCl  
**FW:** 250.2  
**Purity:**  $\geq$ 98%  
**Supplied as:** A crystalline solid  
**Storage:** -20°C  
**Stability:**  $\geq$ 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

Clorprenaline (hydrochloride) is supplied as a crystalline solid. A stock solution may be made by dissolving the clorprenaline (hydrochloride) in the solvent of choice, which should be purged with an inert gas. Clorprenaline (hydrochloride) is soluble in the organic solvent DMSO at a concentration of approximately 1 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of clorprenaline (hydrochloride) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of clorprenaline (hydrochloride) in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

### Description

Clorprenaline is an agonist of the  $\beta_2$ -adrenergic receptor ( $\beta_2$ -AR).<sup>1</sup> Formulations containing clorprenaline have been used to promote growth and improve feed efficiency in livestock.

### Reference

1. Shao, X., Zhang, J., Li, D., *et al.* Preparation of Fe<sub>2</sub>O<sub>3</sub>-clorprenaline/tetraphenylborate nanospheres and their application as ion selective electrode for determination of clorprenaline in pork. *Nanoscale Res. Lett.* **11**(1), 178 (2016).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/04/2020

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM