



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION



## Palmitic Acid MaxSpec® Standard

Item No. 30380

CAS Registry No.: 57-10-3

Formal Name: hexadecanoic acid

Synonyms: C16:0, Cetyllic Acid, Hexadecanoic Acid, NSC 5030

MF: C<sub>16</sub>H<sub>32</sub>O<sub>2</sub>

FW: 256.4

Purity: ≥95%


Supplied as: A solution in chloroform; in a deactivated glass ampule

Concentration: 100 µg/ml (nominal); see certificate of analysis for verified concentration

Storage: -20°C

Stability: ≥5 years; *Stability testing is ongoing to ensure concentration accuracy. The certificate of analysis and product expiry date will be updated upon completion of testing.*

Special Conditions: Store upright and unopened at -20°C. Warm to room temperature prior to opening. Light sensitive.



### Description

Palmitic acid is a long-chain saturated fatty acid that comprises approximately 25 and 65% of human total plasma lipids and saturated fatty acids, respectively.<sup>1,2</sup> Acylation of palmitic acid to proteins facilitates anchoring of membrane-bound proteins to the lipid bilayer and trafficking of intracellular proteins, promotes protein-vesicle interactions, and regulates various G protein-coupled receptor functions.<sup>1</sup> Palmitic acid (200 µM) increases NF-κB p65 levels, matrix metalloproteinase-9 (MMP-9) activity, and production of reactive oxygen species (ROS) in AsPC-1 pancreatic cancer cells, as well as increases migration of AsPC-1 cells.<sup>3</sup> It increases COX-2 levels in RAW 264.7 cells and increases LPS-induced IL-1β levels and caspase-1 activity in isolated mouse peritoneal macrophages.<sup>4,5</sup> Dietary administration of palmitic acid (2.2% w/w for 12 weeks) increases mouse hippocampal β-secretase 1 (BACE1) activity and amyloid β (1-42) (Aβ42; Item No. 20574) levels.<sup>6</sup> It also induces systolic contractile dysfunction in isolated mouse hearts.<sup>7</sup> Red blood cell palmitic acid levels are increased in patients with metabolic syndrome compared to patients without metabolic syndrome and are also increased in the plasma of patients with type 2 diabetes compared to individuals without diabetes.<sup>8,9</sup>

Palmitic acid MaxSpec® standard is a quantitative grade standard of palmitic acid (Item No. 10006627) that has been prepared specifically for mass spectrometry or any application where quantitative reproducibility is required. The solution has been prepared gravimetrically and is supplied in a deactivated glass ampule sealed under argon. The concentration was verified by comparison to an independently prepared calibration standard. This palmitic acid MaxSpec® standard is guaranteed to meet identity, purity, stability, and concentration specifications and is provided with a batch-specific certificate of analysis. Ongoing stability testing is performed to ensure the concentration remains accurate throughout the shelf life of the product.

**Note:** The amount of solution added to the vial is in excess of the listed amount. Therefore, it is necessary to accurately measure volumes for preparation of calibration standards. Follow recommended storage and handling conditions to maintain product quality.

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 08/07/2023

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897  
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM

# PRODUCT INFORMATION



## References

1. Fatima, S., Hu, X., Gong, R.-H., *et al.* Palmitic acid is an intracellular signaling molecule involved in disease development. *Cell. Mol. Life Sci.* **76(13)**, 2547-2557 (2019).
2. Santos, M.J., López-Jurado, M., Llopis, J., *et al.* Influence of dietary supplementation with fish oil on plasma fatty acid composition in coronary heart disease patients. *Ann. Nutr. Metab.* **39(1)**, 52-62 (1995).
3. Binker-Cosen, M.J., Richards, D., Oliver, B., *et al.* Palmitic acid increases invasiveness pancreatic cancer cells AsPC-1 through TLR4/ROS/NF- $\kappa$ B/MMP-9 signaling pathway. *Biochem. Biophys. Res. Commun.* **484(1)**, 152-158 (2017).
4. Lee, J.Y., Sohn, K.H., Rhee, S.H., *et al.* Saturated fatty acids, but not unsaturated fatty acids, induced the expression of cyclooxygenase-2 mediated through Toll-like receptor 4. *J. Biol. Chem.* **276(20)**, 16683-16689 (2001).
5. Karasawa, T., Kawashima, A., Usui-Kawanishi, F., *et al.* Saturated fatty acids undergo intracellular crystallization and activate the NLRP3 inflammasome in macrophages. *Arterioscler. Thromb. Vasc. Biol.* **38(4)**, 744-756 (2018).
6. Marwarha, G., Rostad, S., Lilek, J., *et al.* Palmitate increases  $\beta$ -site A $\beta$ PP-cleavage enzyme 1 activity and amyloid- $\beta$  genesis by evoking endoplasmic reticulum stress and subsequent C/EBP homologous protein activation. *J. Alzheimers Dis.* **57(3)**, 907-925 (2017).
7. Knowles, C.J., Cebova, M., and Pinz, I.M. Palmitate diet-induced loss of cardiac caveolin-3: A novel mechanism for lipid-induced contractile dysfunction. *PLoS One* **8(4)**, e61369 (2013).
8. Yi, L.-Z., He, J., Liang, Y.-Z., *et al.* Plasma fatty acid metabolic profiling and biomarkers of type 2 diabetes mellitus based on GC/MS and PLS-LDA. *FEBS Lett.* **580(30)**, 6837-6845 (2006).
9. Kabagambe, E.K., Tsai, M.Y., Hopkins, P.N., *et al.* Erythrocyte fatty acid composition and the metabolic syndrome: A National Heart, Lung, and Blood Institute GOLDN study. *Clin. Chem.* **54(1)**, 154-162 (2008).

### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete Safety Data Sheet](#), which has been sent via email to your institution.

### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/02/2020

## CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897  
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM