

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



α -D-Glucose-1-phosphate (sodium salt hydrate)

Item No. 30566

CAS Registry No.:	150399-99-8	
Formal Name:	1-(dihydrogen phosphate)α-D-	ОН
Synonym:	glucopyranose tetrahydrate, disodium salt α-D-Glc 1-P	HONOHO
MF:	C ₆ H ₁₁ O ₉ P ● 2Na [4H ₂ O]	
FW:	376.2	HO P OF
Purity:	≥95%	0 0
Supplied as:	A crystalline solid	• 2Na ⁺ [4H ₂ O]
Storage:	-20°C	
Stability:	≥2 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.		

Laboratory Procedures

 α -D-Glucose-1-phosphate (sodium salt hydrate) is supplied as a crystalline solid. Aqueous solutions of α -D-glucose-1-phosphate (sodium salt hydrate) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of α -D-glucose-1-phosphate (sodium salt hydrate) in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

 α -D-Glucose-1-phosphate is an intermediate in glycogen metabolism.^{1,2} It is a precursor in the biosynthesis of UDP-glucose, the glucose donor in glycogen biosynthesis.² α -D-Glucose-1-phosphate can be formed during glycogen breakdown via phosphorolytic cleavage of glycogen by glycogen phosphorylase.¹ It can be converted to glucose-6-phosphate by phosphoglucomutase. α -D-Glucose-1-phosphate is combined with CTP by α -D-glucose-1-phosphate cytidylyltransferase to form CDP-glucose in the first step of CDP-D-tyvelose biosynthesis in S. typhi.³ Bacterial sucrose phosphorylase converts sucrose and phosphate into D-fructose and α -D-glucose-1-phosphate.⁴

References

- 1. Berg, J.M., Tymoczko, J.L., and Stryer, L. Glycogen breakdown requires the interplay of several enzymes. Biochemistry. 5th edition, W H Freeman (2002).
- 2. Berg, J.M., Tymoczko, J.L., and Stryer, L. Glycogen is synthesized and degraded by different pathways. Biochemistry, 5th edition, W H Freeman (2002).
- 3. Koropatkin, N.M. and Holden, H.M. Molecular structure of a-D-glucose-1-phosphate cytidylyltransferase from Salmonella typhi. J. Biol. Chem. 279(42), 44023-44029 (2004).
- 4. Goedl, C., Schwarz, A., Minani, A., et al. Recombinant sucrose phosphorylase from Leuconostoc mesenteroides: Characterization, kinetic studies of transglucosylation, and application of immobilised enzyme for production of a-D-glucose 1-phosphate. J. Biotechnol. 129(1), 77-86 (2007).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/08/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM