

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



ONC212

Item No. 30931

CAS Registry No.:	1807861-48-8	
Formal Name:	2,4,6,7,8,9-hexahydro-7-	
	(phenylmethyl)-4-[[4-(trifluoromethyl) phenyl]methyl]-imidazo[1,2-a] pyrido[3,4-e]pyrimidin-5(1H)-one	
MF:	$C_{24}H_{23}F_{3}N_{4}O$	
FW:	440.5	
Purity:	≥98%	v v v H v v
UV/Vis.:	λ _{max} : 232 nm	0
Supplied as:	A crystalline solid	
Storage:	-20°C	
Stability:	≥2 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.		

Laboratory Procedures

ONC212 is supplied as a crystalline solid. A stock solution may be made by dissolving the ONC212 in the solvent of choice, which should be purged with an inert gas. ONC212 is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of ONC212 in these solvents is approximately 1 mg/ml.

ONC212 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, ONC212 should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. ONC212 has a solubility of approximately 0.25 mg/ml in a 1:3 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

ONC212 is an agonist of the orphan G protein-coupled receptor GPR132 (EC $_{50}$ = ~400 nM in a PathHunter[®] β-arrestin assay).¹ It inhibits cell growth in a panel of 62 human leukemia cell lines, including acute myeloid leukemia (AML), acute lymphoblastic leukemia (ALL), chronic myelogenous leukemia (CML), and hairy cell leukemias, with GI₅₀ values ranging from less than 78 to 456 nM. ONC212 also reduces cell growth in a panel of 58 lymphoma cell lines (GI₅₀s = <78-261 nM). In vivo, ONC212 (50 mg/kg) reduces tumor volume in HT-29 and HCT116 mouse xenograft models.²

References

- 1. Prabhu, V.V., Madhukar, N., Tarapore, R., et al. Abstract 1155: Potent anti-cancer effects of selective GPR132/G2A agonist imipridone ONC212 in leukemia and lymphoma. Cancer Res. 77(13), (2017).
- 2. Wagner, J., Kline, C.L., Ralff, M.D., et al. Preclinical evaluation of the imipridone family, analogs of clinical stage anti-cancer small molecule ONC201, reveals potent anti-cancer effects of ONC212. Cell Cycle 16(19), 1790-1799 (2017).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 08/24/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM