



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



4-Hydroxyestradiol

Item No. 31508

CAS Registry No.: 5976-61-4
Formal Name: (17 β)-estra-1,3,5(10)-triene-3,4,17-triol
Synonym: 4-Hydroxy-17 β -estradiol
MF: C₁₈H₂₄O₃
FW: 288.4
Purity: \geq 98%
Supplied as: A crystalline solid
Storage: -20°C
Stability: \geq 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

4-Hydroxyestradiol is supplied as a crystalline solid. A stock solution may be made by dissolving the 4-hydroxyestradiol in the solvent of choice, which should be purged with an inert gas. 4-Hydroxyestradiol is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of 4-hydroxyestradiol in ethanol and DMSO is approximately 25 mg/ml and approximately 20 mg/ml in DMF.

4-Hydroxyestradiol is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, 4-hydroxyestradiol should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. 4-Hydroxyestradiol has a solubility of approximately 0.25 mg/ml in a 1:3 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

4-Hydroxyestradiol is a catechol estrogen and minor metabolite of estrogen.^{1,2} 4-Hydroxyestradiol is formed from estrogen by the cytochrome P450 (CYP) isoform CYP1B1.² Production of 4-hydroxyestradiol is increased in microsomes prepared from human mammary adenocarcinoma and fibroadenoma microsomes compared with microsomes prepared from non-cancerous tissues.¹ 4-Hydroxyestradiol (25 mg/animal) induces renal tumor formation in male hamsters.³

References

1. Liehr, J.C. and Ricci, M.J. 4-Hydroxylation of estrogens as marker of human mammary tumors. *Proc. Natl. Acad. Sci. USA* **93**(8), 3294-3296 (1996).
2. Yager, J.D. and Liehr, J.G. Molecular mechanisms of estrogen carcinogenesis. *Annu. Rev. Pharmacol. Toxicol.* **36**, 203-232 (1996).
3. Liehr, J.G., Fang, W.F., Sirbasku, D.A., et al. Carcinogenicity of catechol estrogens in Syrian hamsters. *J. Steroid Biochem.* **24**(1), 353-356 (1986).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/18/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM