



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



Prostaglandin F_{2α}-d₄ Item No. 316010

CAS Registry No.: 34210-11-2
Formal Name: 9α,11α,15S-trihydroxy-prosta-5Z,13E-dien-1-oic-3,3,4,4-d₄ acid
Synonyms: Dinoprost-d₄, PGF_{2α}-d₄
MF: C₂₀H₃₀D₄O₅
FW: 358.5
Chemical Purity: ≥98%
Deuterium Incorporation: ≥99% deuterated forms (d₁-d₄); ≤1% d₀
Supplied as: A solution in methyl acetate
Storage: -20°C
Stability: As supplied, 1 year from the QC date provided on the Certificate of Analysis, when stored properly



Laboratory Procedures

Prostaglandin F_{2α}-d₄ (PGF_{2α}-d₄) contains four deuterium atoms at the 3, 3', 4, and 4' positions. It is intended for use as an internal standard for the quantification of PGF_{2α} (Item No. 16010) by GC- or LC-mass spectrometry (MS).

PGF_{2α}-d₄ is supplied as a solution in methyl acetate. To change the solvent, simply evaporate the methyl acetate under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of PGF_{2α}-d₄ in these solvents is approximately 100 mg/ml.

PGF_{2α}-d₄ is used as an internal standard for the quantification of PGF_{2α} by stable isotope dilution MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the weight indicated on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard PGF_{2α} by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Description

PGF_{2α} is a widely distributed PG occurring in many species.¹⁻³ It causes contraction of vascular, bronchial, intestinal, and myometrial smooth muscle, and also exhibits potent luteolytic activity.¹ PGF_{2α} exerts its receptor mediated physiological activity at 50-100 nM.¹ Maximal ovine myometrial contraction can be achieved at 125 nM PGF_{2α} *in vitro*.⁴

References

1. Samuelsson, B., Goldyne, M., Granström, E., *et al. Annu. Rev. Biochem.* **47**, 997-1029 (1978).
2. Speroff, L. and Ramwell, P.W. *Am. J. Obstet. Gynecol.* **107**, 1111-1130 (1970).
3. Watanabe, K., Iguchi, Y., Iguchi, S., *et al. Proc. Natl. Acad. Sci. USA* **83**, 1583-1587 (1986).
4. Crankshaw, D.J. and Gaspar, V. *J. Reprod. Fertil.* **103**, 55-61 (1995).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 04/07/2016

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM