

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Dermorphin (acetate)

Item No. 31607

Formal Name:	(S)-1-(L-tyrosyl-D-alanyl-L- phenylalanylglycyl-L-tyrosyl)- N-((S)-1-amino-3-hydroxy-1- oxopropan-2-yl)pyrrolidine-2- carboxamide, acetate	
MF: FW:	$C_{40}H_{50}N_8O_{10} \bullet XC_2H_4O_2$ 802.9	
Purity:	≥95%	
UV/Vis.:	λ _{max} : 224 nm	
Supplied as:	A crystalline solid	H ₂ N
Storage:	-20°C	• XCH3CO2H
Stability:	≥2 years	Хон. ₃ оо ₂ `ОН

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Dermorphin (acetate) is supplied as a crystalline solid. A stock solution may be made by dissolving the dermorphin (acetate) in the solvent of choice, which should be purged with an inert gas. Dermorphin (acetate) is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of dermorphin (acetate) in these solvents is approximately 30 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of dermorphin (acetate) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of dermorphin (acetate) in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Dermorphin is an opioid peptide originally isolated from the skin of South American frogs (Ph. sauvagei).¹ It binds to μ -opioid receptors (K_i = 0.54 nM) and is selective for μ - over δ -opioid receptors in radioligand binding assays (K₁ = 929 nM). In vivo, dermorphin (1.9 µmol/kg, i.p.) inhibits noxious stimuli-induced neuronal firing in the nucleus lateralis anterior and ventrobasal complex in rats.² It inhibits the electrical stimulation-induced C fiber response in rat dorsal horn nociceptive neurons. Dermorphin (10-120 pmol/animal, i.c.v.) delays gastric emptying, inhibits gastric acid secretion, and slows colonic motility in pylorus-ligated rats.

References

- 1. Usenko, A.B., Emel'yanova, T.G., and Mlasoedov, N.F. Dermorphins are natural opioids with an unique primary structure that determines their biological specificity. Biol. Bull. Russ. Acad. Sci. 29, 154-164 (2002).
- 2. Melchiorri, P. and Negri, L. The dermorphin peptide family. Gen. Pharmacol. 27(7), 1099-1107 (1996).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/23/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM