



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION



## LIF (human, recombinant)

Item No. 32066

### Overview and Properties

**Synonyms:** Differentiation Inhibitory Factor, HILDA, Human Interleukin in DA Cells, Leukemia Inhibitor Factor, LIF Interleukin 6 Family Cytokine, Melanoma-derived LPL Inhibitor, MLPLI

**Source:** Active recombinant human LIF expressed in HEK293 cells

**Amino Acids:** 23-202

**Uniprot No.:** P15018

**Molecular Weight:** 19.7 kDa

**Storage:** -80°C (as supplied)

**Stability:** ≥1 year

**Purity:** ≥95% estimated by SDS-PAGE


**Supplied in:** Lyophilized from sterile PBS, pH 7.4

**Endotoxin Testing:** <1.0 EU/μg, determined by the LAL endotoxin assay


**Bioactivity:** See figures for details

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Image(s)



**SDS-PAGE Analysis of LIF.** This protein has a calculated molecular weight of 19.7 kDa. It has an apparent molecular weight of approximately 35.4 kDa by SDS-PAGE under reducing conditions due to glycosylation.



**Ability of LIF to inhibit the proliferation of M1 mouse myeloid leukemia cells.** Measured by its ability to inhibit the proliferation of M1 mouse myeloid leukemia cells. The EC<sub>50</sub> for this effect is typically 0.2-0.8 ng/mL.

**WARNING**  
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

**SAFETY DATA**  
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

**WARRANTY AND LIMITATION OF REMEDY**  
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/22/2020

**CAYMAN CHEMICAL**  
1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA  
PHONE: [800] 364-9897  
[734] 971-3335  
FAX: [734] 971-3640  
CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM

# PRODUCT INFORMATION



## Description

---

Leukemia inhibitory factor (LIF) is a cytokine and member of the IL-6 family with roles in cell growth and differentiation, bone metabolism, and inflammation.<sup>1</sup> Mature LIF is a 180-amino acid peptide arranged as a four-helix bundle in an up-up-down-down configuration that is synthesized as a 202-amino acid precursor which is post-translationally processed to remove a 22-amino acid signaling sequence from its N-terminus. It is ubiquitously expressed and this expression is commonly upregulated under inflammatory conditions.<sup>2</sup> LIF binds to its receptor LIFR $\beta$  and recruits the IL-6 family common receptor gp130 to form a heterodimer and induce intracellular signaling via the JAK-STAT pathway.<sup>1,3</sup> Exogenous administration of LIF reduces the rate of deterioration in grip strength and slows progression of the forelimb deformity in the wobbler mouse model of amyotrophic lateral sclerosis (ALS).<sup>4</sup> Intracerebroventricular administration of a recombinant adeno-associated viral vector encoding LIF reduces food intake and body weight gain in rats.<sup>5</sup> Knockdown of *Lif* induces infertility in female mice.<sup>1</sup> Cayman's LIF (human, recombinant) protein can be used for cell-based assay applications. This protein consists of 180 amino acids, has a calculated molecular weight of 19.7 kDa, and a predicted N-terminus of Ser23 after signal peptide cleavage. By SDS-PAGE, under reducing conditions, the apparent molecular mass of the protein is 35.4 kDa due to glycosylation.

## References

---

1. Nicola, N.A. and Babon, J.J. Leukemia inhibitory factor (LIF). *Cytokine Growth Factor Rev.* **26(5)**, 533-544 (2015).
2. Hisaka, T., Desmoulière, A., Taupin, J.-L., *et al.* Expression of leukemia inhibitory factor (LIF) and its receptor gp190 in human liver and in cultured human liver myofibroblasts. Cloning of new isoforms of LIF mRNA. *Comp. Hepatol.* **3(10)**, (2004).
3. Onishi, K. and Zandstra, P.W. LIF signaling in stem cells and development. *Development* **142(13)**, 2230-2236 (2015).
4. Kurek, J.B., Radford, A.J., Crump, D.E., *et al.* LIF (AM424), a promising growth factor for the treatment of ALS. *J. Neurol. Sci.* **160(Suppl. 1)**, S106-S113 (1998).
5. Pasquin, S., Sharma, M., and Gauchat, J.-F. Cytokines of the LIF/CNTF family and metabolism. *Cytokine* **82**, 122-124 (2016).

CAYMAN CHEMICAL  
1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA  
PHONE: [800] 364-9897  
[734] 971-3335  
FAX: [734] 971-3640  
CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM