



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



Soluble TNF- α (mouse, recombinant)


Item No. 32069

Overview and Properties


Synonyms:	DIF, Differentiation-inducing Factor, TNFA, TNFSF2, Tumor Necrosis Factor- α
Source:	Active recombinant mouse TNF- α expressed in <i>E. coli</i>
Amino Acids:	80-235
Uniprot No.:	P06804
Molecular Weight:	17 kDa
Storage:	-80°C (as supplied)
Stability:	≥ 1 year
Purity:	$\geq 98\%$ estimated by SDS-PAGE
Supplied in:	Lyophilized from sterile PBS, pH 7.4, with 5% trehalose, 5% mannitol, and 0.01% Tween 80
Bioactivity:	See figures for details

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Images



SDS-PAGE Analysis of Soluble TNF- α .



L929 Cytotoxicity Assay. TNF- α activity is measured in a cytotoxicity assay using L929 mouse fibrosarcoma cells in the presence of the metabolic inhibitor actinomycin D. The EC_{50} value for this effect is typically 3-30 μ g/ml.

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/01/2023

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Description

TNF- α is a cytokine and a member of the TNF/TNF receptor (TNFR) cytokine superfamily.¹ TNF- α is produced as a 233-amino acid transmembrane precursor protein from which mature, soluble TNF- α is formed by proteolysis.² Soluble TNF- α is a 157-amino acid polypeptide, cleaved from the precursor protein on the extracellular side of the membrane, that forms bell-shaped homotrimers with the C-termini at the base, each containing three receptor interaction sites.³ It is primarily produced by activated macrophages but can also be produced by a variety of other cells, such as T cells, natural killer cells, and osteoblasts.^{3,4} TNF- α binds to and activates its receptors, TNFR1 and TNFR2, which are associated with intracellular protein complexes that activate caspases to induce cell death, induce p38 MAPK signaling, and initiate NF- κ B or AP-1-mediated transcription of immune and inflammatory mediators.⁵ TNF- α promotes inflammation partly by inducing endothelial cells to express adhesion molecules, COX enzymes, and pro-coagulant factors.⁴ Exogenous TNF- α induces death of cancer cells *in vitro*, as well as disrupts tumor vascularization and induces necrosis *in vivo*, but it has tumor promoting properties when produced in the cancer microenvironment.^{1,6} In contrast, it plays a role in resistance to infection, with mice lacking *Tnf* having an increased susceptibility to certain microbial infections but lacking resistance to leishmania.⁵ *Tnf* knockout mice are also resistant to certain types of cancer, including chemically induced skin carcinogenesis.¹ TNF- α increases lung metastases in a mouse model of fibrosarcoma, an effect that can be reduced by an anti-TNF- α antibody. Mice overexpressing *Tnf* develop an arthritis similar to rheumatoid arthritis in humans.⁷ TNF- α is produced in the inflamed tissues of patients with inflammatory diseases such as rheumatoid arthritis, and neutralizing antibodies to TNF- α reduce the levels of TNF- α *in vitro* and in mouse models of the disease.⁴ Cayman's Soluble TNF- α (mouse, recombinant) protein can be used for cell-based assay applications. This protein consists of 157 amino acids and has a calculated molecular weight of 17 kDa.

References

1. Balkwill, F. TNF- α in promotion and progression of cancer. *Cancer Metastasis Rev.* **25(3)**, 409-416 (2006).
2. Kriegler, M., Perez, C., DeFray, K., *et al.* A novel form of TNF/cachectin is a cell surface cytotoxic transmembrane protein: Ramifications for the complex physiology of TNF. *Cell* **53(1)**, 45-53 (1988).
3. Tang, P., Hung, M., and Klostergaard, J. Human pro-tumor necrosis factor is a homotrimer. *Biochemistry* **35(25)**, 8216-8225 (1996).
4. Bradley, J.R. TNF-mediated inflammatory disease. *J. Pathol.* **214(2)**, 149-160 (2008).
5. Idriss, H.T. and Naismith, J.H. TNF α and the TNF receptor superfamily: Structure-function relationship(s). *Microsc. Res. Tech.* **50(3)**, 184-195 (2000).
6. Josephs, S.F., Ichim, T.E., Prince, S.M., *et al.* Unleashing endogenous TNF-alpha as a cancer immunotherapeutic. *J. Transl. Med.* **16(1)**, 242 (2018).
7. Li, P. and Schwarz, E.M. The TNF- α transgenic mouse model of inflammatory arthritis. *Springer Semin. Immunopathol.* **25(1)**, 19-33 (2003).

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM