



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



15-(6-nitroxyhexanoyl)-17-phenyl trinor Prostaglandin F_{2α}

Item No. 32791

CAS Registry No.: 1194396-71-8

Formal Name: 6-(nitrooxy)-hexanoic acid, (1S,2E)-3-
[[1R,2R,3S,5R)-2-[(2Z)-7-(ethylamino)-7-oxo-2-
hepten-1-yl]-3,5-dihydroxycyclopentyl]-1-(2-
phenylethyl)-2-propen-1-yl ester

Synonyms: 15-(6-nitroxyhexanoyl)-17-phenyl trinor PGF_{2α},
15-(6-nitroxyhexanoyl)-Bimatoprost,
NCX 470

MF: C₃₁H₄₆N₂O₈


FW: 574.7

Purity: ≥98%

Supplied as: A solution in ethanol

Storage: -20°C

Stability: ≥1 year



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

15-(6-nitroxyhexanoyl)-17-phenyl trinor Prostaglandin F_{2α} (15-(6-nitroxyhexanoyl)-17-phenyl trinor PGF_{2α}) is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO purged with an inert gas can be used. The solubility of 15-(6-nitroxyhexanoyl)-17-phenyl trinor PGF_{2α} in DMSO is approximately 10 mg/ml.

Description

15-(6-nitroxyhexanoyl)-17-phenyl trinor PGF_{2α} is a nitric oxide-donating derivative of 17-phenyl trinor PGF_{2α} (Item No. 16810).¹ It increases cGMP levels in rabbit aqueous humor and iris ciliary body when topically administered at a concentration of 0.042%. Topical administration of 15-(6-nitroxyhexanoyl)-17-phenyl trinor PGF_{2α} (0.14%) reduces intraocular pressure (IOP) in a rabbit model of hypertonic saline-induced transient ocular hypertension. It also reduces IOP in a cynomolgus monkey model of laser-induced ocular hypertension when administered topically at a concentration of 0.042%.

Reference

1. Impagnatiello, F., Toris, C.B., Batugo, M., *et al.* Intraocular pressure-lowering activity of NCX 470, a novel nitric oxide-donating bimatoprost in preclinical models. *Invest. Ophthalmol. Vis. Sci.* **56(11)**, 6558-6564 (2015).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/06/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM