



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Benastatin C

Item No. 32894

CAS Registry No.: 150151-88-5
Formal Name: 1,7,9,11-tetrahydroxy-13,13-dimethyl-3-pentyl-benzo[a]naphthacen-8(13H)-one
MF: C₂₉H₂₈O₅
FW: 456.5
Purity: ≥90%
Supplied as: A powder
Storage: -20°C
Stability: ≥2 years
Item Origin: Bacterium/*Streptomyces* sp. MI384-DF12



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Benastatin C is supplied as a powder. A stock solution may be made by dissolving the benastatin C in the solvent of choice, which should be purged with an inert gas. Benastatin C is soluble in acetone, ethyl acetate, methanol, and DMSO.

Description

Benastatin C is a polyketide synthase-derived benastatin that has been found in *Streptomyces* and has diverse biological activities.^{1,2} It inhibits glutathione S-transferase (GST; IC₅₀ = 24 µg/ml for the rat liver enzyme).² Benastatin C also inhibits the esterase activity of isolated porcine pancreatic lipase (IC₅₀ = 10 µg/ml). It increases LPS- or concanavalin A-induced blastogenesis of isolated mouse spleen lymphocytes in a concentration-dependent manner.

References

- Xu, Z., Schenk, A., and Hertweck, C. Molecular analysis of the benastatin biosynthetic pathway and genetic engineering of altered fatty acid-polyketide hybrids. *J. Am. Chem. Soc.* **129(18)**, 6022-6030 (2007).
- Aoyama, T., Kojima, F., Yamazaki, T., et al. Benastatins C and D, new inhibitors of glutathione S-transferase, produced by *Streptomyces* sp. MI384-DF12. Production, isolation, structure determination and biological activities. *J. Antibiot. (Tokyo)* **46(5)**, 712-718 (1993).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/24/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM