



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



17 α -Hydroxyprogesterone

Item No. 33154

CAS Registry No.: 68-96-2
Formal Name: 17-hydroxy-pregn-4-ene-3,20-dione
Synonyms: NSC 15468, 17-OHP, 17-hydroxy Progesterone
MF: C₂₁H₃₀O₃
FW: 330.5
Purity: \geq 98%
UV/Vis.: λ_{max} : 241 nm
Supplied as: A crystalline solid
Storage: -20°C
Stability: \geq 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

17 α -Hydroxyprogesterone (17-OHP) is supplied as a crystalline solid. A stock solution may be made by dissolving the 17-OHP in the solvent of choice, which should be purged with an inert gas. 17-OHP is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of 17-OHP in these solvents is approximately 30 mg/ml.

Description

17-OHP is a steroid and precursor in the synthesis of adrenal androgens.^{1,2} It binds to the glucocorticoid receptor in a cell-free assay (IC₅₀ = 82 nM for the human receptor) and induces glucocorticoid receptor transactivation in HEK293 cells (EC₅₀ = 1.2 μ M).² 17-OHP (0.005-1 μ M) inhibits aldosterone-induced transactivation of the mineralocorticoid receptor in cell-based assays.¹ Serum levels of 17-OHP are increased in patients with congenital adrenal hyperplasia.^{1,2}

References

1. Mooij, C.F., Parajes, S., Pijnenburg-Kleizen, K.J., *et al.* Influence of 17-hydroxyprogesterone, progesterone and sex steroids on mineralocorticoid receptor transactivation in congenital adrenal hyperplasia. *Horm. Res. Paediatr.* **83(6)**, 414-421 (2015).
2. Pijnenburg-Kleizen, K.J., Engels, M., Mooij, C.F., *et al.* Adrenal steroid metabolites accumulating in congenital adrenal hyperplasia lead to transactivation of the glucocorticoid receptor. *Endocrinology* **156(10)**, 3504-3510 (2015).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/03/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM