

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



# Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



СООН

# (±)4-HDHA

Item No. 33200

CAS Registry No.: 90906-40-4

Formal Name: (±)4-hydroxy-5E,7Z,10Z,13Z,16Z,19Z-

docosahexaenoic acid

Synonyms: 4-hydroxy Docosahexaenoic Acid,

(±)4-HDoHE

MF:  $C_{22}H_{32}O_3$ FW: 344.5 **Purity:** ≥95%

UV/Vis.:  $\lambda_{max}$ : 236 nm  $\epsilon$ : 20,000 Supplied as: A solution in ethanol

-20°C Storage:

Stability: As supplied, 2 years from the QC date provided on the Certificate of Analysis, when

stored properly

#### **Laboratory Procedures**

(±)4-HDHA is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO and dimethyl formamide purged with an inert gas can be used. (±)4-HDHA is miscible in these solvents.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of (±)4-HDHA is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of (±)4-HDHA in PBS (pH 7.2) is approximately 0.8 mg/ml. We do not recommend storing the aqueous solution for more than one day.

## Description

(±)4-HDHA is an autoxidation product of docosahexaenoic acid (DHA) in vitro. 1,2 It is also produced from incubations of DHA in rat liver, brain, and intestinal microsomes.<sup>3-5</sup> Enzymatic transformation of DHA by RBL-1 cells also produces 4-HDHA.<sup>6</sup> However, the enzymatic product is most likely the S-isomer. (±)4-HDHA is a potential marker of oxidative stress in brain and retina where DHA is an abundant polyunsaturated fatty acid.

## References

- 1. VanRollins, M. and Murphy, R.C. J. Lipid Res. 25, 507-517 (1984).
- 2. Reynaud, D., Thickitt, C.P., and Pace-Asciak, C.R. Anal. Biochem. 214, 165-170 (1993).
- 3. VanRollins, M., Baker, R.C., Sprecher, H., et al. J. Biol. Chem. 259, 5776-5783 (1984).
- 4. Yamane, M., Abe, A., and Yamane, S. Journal of Chromatography B 652, 123-136 (1994).
- 5. Kim, H.Y., Karanian, J.W., Shingu, T., et al. Prostaglandins 40, 473-491 (1990).
- Corey, E.J., Shih, C., and Cashman, J.R. Proc. Natl. Acad. Sci. USA 80, 3581-3584 (1983).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

al should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution

## WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 07/28/2016

#### **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA **PHONE:** [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM