

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Deferitrin

Item No. 33328

CAS Registry No.:	239101-33-8	
Formal Name:	(4S)-2-(2,4-dihydroxyphenyl)-4,5-dihydro-4-	
	methyl-4-thiazolecarboxylic acid	0
Synonyms:	GT-56-252, (S)-4'-Hydroxydesazadesferrithiocin	ş
MF:	C ₁₁ H ₁₁ NO ₄ S	ОН
FW:	253.3	N *
Purity:	≥98%	
UV/Vis.:	λ _{max} : 214, 231, 274, 309 nm	
Supplied as:	A solid	но. 🗸 он
Storage:	-20°C	
Stability:	≥4 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.		

Laboratory Procedures

Deferitrin is supplied as a solid. A stock solution may be made by dissolving the deferitrin in the solvent of choice, which should be purged with an inert gas. Deferitrin is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of deferitrin in these solvents is approximately 12 and 10 mg/ml, respectively. Deferitrin is slightly soluble in ethanol.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of deferitrin can be prepared by directly dissolving the solid in aqueous buffers. The solubility of deferitrin in PBS (pH 7.2) is approximately 0.2 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Deferitrin is an iron chelator and a derivative of desferrithiocin that binds iron in a 2:1 (ligand:iron) ratio.^{1,2} It inhibits the proliferation of L1210 murine leukemia cells (IC₅₀ = 17 μ M).³ Deferitrin (150 μ mol/kg) induces iron excretion in iron-overloaded C. apella monkeys and beagle dogs with an iron clearing efficiency (ICE) of 14.5 and 9.3%, respectively.² It reduces fecal and urine iron levels by 607 and 357 μ g/kg in *C. apella* monkeys and beagle dogs, respectively.

References

- 1. Bergeron, R.J., Wiegand, J., Bharti, N., et al. Desferrithiocin analogue iron chelators: Iron clearing efficiency, tissue distribution, and renal toxicity. Biometals 24(2), 239-258 (2011).
- 2. Bergeron, R.J., Wiegand, J., Weimar, W.R., et al. Comparison of iron chelator efficacy in iron-overloaded beagle dogs and monkeys (Cebus apella). Comp. Med. 54(6), 664-672 (2004).
- 3. Bergeron, R.J., Wiegand, J., Weimar, W.R., et al. Iron chelation promoted by desazadesferrithiocin analogs: An enantioselective barrier. Chirality 15(7), 593-599 (2003).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/27/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM