

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

# **PRODUCT** INFORMATION



### Chlormethiazole (hydrochloride)

Item No. 33437

| CAS Registry No.: | 6001-74-7                                |                        |   |
|-------------------|--|------------------------|---|
| Formal Name:      | 5-(2-chloroethyl)-4-methyl-              |                        |   |
|                   | thiazole, monohydrochloride              |                        |   |
| Synonym:          | Clomethiazole                            | ,N                     |   |
| MF:               | C <sub>6</sub> H <sub>8</sub> CINS ● HCI |                        |   |
| FW:               | 198.1                                    |                        |   |
| Purity:           | ≥98%                                     | 3. <u> </u>            | 1 |
| UV/Vis.:          | λ <sub>max</sub> : 213, 250 nm           | • HCI                  |   |
| Supplied as:      | A crystalline solid                      |                        |   |
| Storage:          | -20°C                                    |                        |   |
| Stability:        | ≥2 years                                 |                        |   |
| 1 ( )             |  | с <u>і с і і і і і</u> |   |

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

#### Laboratory Procedures

Chlormethiazole (hydrochloride) is supplied as a crystalline solid. A stock solution may be made by dissolving the chlormethiazole (hydrochloride) in the solvent of choice, which should be purged with an inert gas. Chlormethiazole (hydrochloride) is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of chlormethiazole (hydrochloride) in ethanol is approximately 20 mg/ml and approximately 30 mg/ml in DMSO and DMF.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of chlormethiazole (hydrochloride) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of chlormethiazole (hydrochloride) in PBS (pH 7.2) is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

#### Description

Chlormethiazole is a potentiator of GABA<sub>A</sub> receptors.<sup>1</sup> It potentiates GABA-induced currents in L-M(TK-) cells expressing  $\alpha_1\beta_1\gamma_2$  or  $\alpha_1\beta_2\gamma_2$  subunit-containing GABA receptors with EC<sub>50</sub> values of 0.8 and 3.4  $\mu$ M, respectively. Chlormethiazole protects against seizures induced by maximal electroshock (MES) in mice  $(ED_{50} = 130.8 \text{ mg/kg})^2$  It reduces infarct volume in a rat model of focal ischemia induced by permanent middle cerebral artery occlusion (MCAO).<sup>3</sup> Formulations containing chlormethiazole have been used in the treatment of acute alcohol withdrawal.

#### References

- 1. Nelson, R.M., Green, A.R., and Hainsworth, A.H. Electrophysiological actions of γ-aminobutyric acid and clomethiazole on recombinant GABA, receptors. Eur. J. Pharmacol. 452(3), 255-262 (2002).
- 2. Pilip, S., Urbańska, E.M., Swiader, M., et al. Anticonvulsant action of chlormethiazole is prevented by subconvulsive amounts of strychnine and aminophylline but not by bicuculline and picrotoxin. Pol. J. Pharmacol. 52(4), 267-273 (2000).
- 3. Sydserff, S.G., Cross, A.J., and Green, A.R. The neuroprotective effect of chlormethiazole on ischaemic neuronal damage following permanent middle cerebral artery ischaemia in the rat. Neurodegeneration 4(3), 323-328 (1995).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/01/2021

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM