



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



Roxatidine-d₁₀ (hemioxalate)

Item No. 34007

Formal Name: 2-hydroxy-N-[3-[3-(1-piperidinyl-2,2,3,3,4,4,5,5,6,6-d₁₀-methyl)phenoxy]propyl]-acetamide, ethanedioate (2:1)

MF: C₁₇H₁₆D₁₀N₂O₃ • 1/2C₂H₂O₄

FW: 361.5


Chemical Purity: ≥95% (Roxatidine)

Deuterium Incorporation: ≥99% deuterated forms (d₁-d₁₀); ≤1% d₀

Supplied as: A solid

Storage: -20°C

Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Roxatidine-d₁₀ (hemioxalate) is intended for use as an internal standard for the quantification of roxatidine by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Roxatidine-d₁₀ (hemioxalate) is supplied as a solid. A stock solution may be made by dissolving the roxatidine-d₁₀ (hemioxalate) in the solvent of choice, which should be purged with an inert gas. Roxatidine-d₁₀ (hemioxalate) is soluble in DMSO.

Description

Roxatidine is a histamine H₂ receptor antagonist and major active metabolite of roxatidine acetate (Item No. 21248).¹ Roxatidine reduces histamine-induced adenylate cyclase production in guinea pig parietal cells (IC₅₀ = 0.8 μM). It inhibits histamine-induced hydrogen ion accumulation in the same cells (pA₂ = 7.03). Roxatidine (200 mg/kg) reduces small intestinal lesion area in a rat model of gastric mucosal injury induced by indomethacin (Item No. 70270).²

References

1. Sewing, K.-F., Beil, W., and Hannemann, H. Comparative pharmacology of histamine H₂-receptor antagonists. *Drugs* **35(Suppl. 3)**, 25-29 (1988).
2. Umegaki, E., Yoda, Y., Tokioka, S., et al. Protective effect of roxatidine against indomethacin-induced small intestinal mucosal injury in rats. *J. Gastroenterol. Hepatol.* **25(Suppl. 1)**, S35-S40 (2010).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/13/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM