



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## CDPPB

Item No. 34132

**CAS Registry No.:** 781652-57-1  
**Formal Name:** 3-cyano-N-(1,3-diphenyl-1H-pyrazol-5-yl)-benzamide  
**MF:** C<sub>23</sub>H<sub>16</sub>N<sub>4</sub>O  
**FW:** 364.4  
**Purity:** ≥98%  
**UV/Vis.:** λ<sub>max</sub>: 264 nm  
**Supplied as:** A solid  
**Storage:** -20°C  
**Stability:** ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

CDPPB is supplied as a solid. A stock solution may be made by dissolving the CDPPB in the solvent of choice, which should be purged with an inert gas. CDPPB is soluble in organic solvents such as DMSO and dimethyl formamide (DMF). The solubility of CDPPB in DMSO and DMF is approximately 25 and 20 mg/ml, respectively.

CDPPB is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, CDPPB should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. CDPPB has a solubility of approximately 0.3 mg/ml in a 1:2 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

### Description

CDPPB is a positive allosteric modulator (PAM) of metabotropic glutamate receptor 5 (mGluR5).<sup>1</sup> It potentiates glutamate-induced increases in intracellular calcium in CHO cells expressing mGluR5 (EC<sub>50</sub>s = 10 and 20 nM for the human and rat receptor, respectively). CDPPB (30 mg/kg, s.c.) inhibits amphetamine-induced disruptions in prepulse inhibition (PPI) of the acoustic startle response in rats.

### Reference

1. Lindsley, C.W., Wisnoski, D.D., Leister, W.H., *et al.* Discovery of positive allosteric modulators for the metabotropic glutamate receptor subtype 5 from a series of N-(1,3-diphenyl-1H-pyrazol-5-yl)benzamides that potentiate receptor function in vivo. *J. Med. Chem.* **47**(24), 5825-5828 (2004).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/15/2021

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM