

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Maltose (hydrate)

Item No. 34239

CAS Registry No.: 6363-53-7

Formal Name: 4-O-α-D-glucopyranosyl-D-glucose,

monohydrate

Synonym: D-(+)-Maltose MF: C₁₂H₂₂O₁₁ • H₂O

360.3 FW: ≥90% **Purity:** Supplied as: A solid Storage: -20°C Stability: ≥2 years

ОН

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Maltose (hydrate) is supplied as a solid. A stock solution may be made by dissolving the maltose (hydrate) in the solvent of choice, which should be purged with an inert gas. Maltose (hydrate) is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of maltose (hydrate) in these solvents is approximately 15 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of maltose (hydrate) can be prepared by directly dissolving the solid in aqueous buffers. The solubility of maltose (hydrate) in PBS (pH 7.2) is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Maltose is a disaccharide found in plants and bacteria. 1,2 It is a degradation product of starch in plants that accumulates in a circadian rhythm-, day length-, and temperature-dependent manner. Maltose is formed from starch by the action of β -amylase and is a source of glucose in bacteria and mammals.^{2,3}

References

- 1. Lu, Y. and Sharkey, T.D. The importance of maltose in transitory starch breakdown. Plant Cell Environ. **29(3)**, 353-366 (2006).
- 2. Boos, W. and Böhm, A. Learning new tricks from an old dog: MalT of the Escherichia coli maltose system is part of a complex regulatory network. Trends Genet. 16(9), 404-409 (2000).
- 3. Tester, R.F., Karkalas, J., and Qi, X. Starch structure and digestibility enzyme-substrate relationship. World Poultry Sci. J. 60(2), 186-195 (2004).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 06/22/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM