

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Tropinone

Item No. 34287

| CAS Registry No.: | 532-24-1 |
|--|---|
| Formal Name: | 8-methyl-8-azabicyclo[3.2.1]octan-3-one |
| Synonyms: | NSC 118012, Tropanone |
| MF: | C ₈ H ₁₃ NO |
| FW: | 139.2 |
| Purity: | ≥98% |
| Supplied as: | A solid |
| Storage: | -20°C |
| Stability: | ≥2 years |
| Item Origin: | Synthetic |
| Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis. | |

Laboratory Procedures

Tropinone is supplied as a solid. A stock solution may be made by dissolving the tropinone in the solvent of choice, which should be purged with an inert gas. Tropinone is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of tropinone in these solvents is approximately 30 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of tropinone can be prepared by directly dissolving the solid in aqueous buffers. The solubility of tropinone in PBS (pH 7.2) is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Tropinone is a polyketide synthase-derived tropane that has been found in A. belladonna.¹⁻³ It is a central intermediate in the biosynthesis of various tropane alkaloids, including (-)-hyoscyamine (Item No. 25644), (+)-hyoscyamine, scopolamine, atropine (Item No. 12008), and cocaine.

References

- 1. Bedewitz, M.A., Jones, A.D., D'Auria, J.C., et al. Tropinone synthesis via an atypical polyketide synthase and P450-mediated cyclization. Nat. Commun. 9(1), 5281 (2018).
- 2. Piechowska, K., Mizerska-Kowalska, M., Zdzisińska, B., et al. Tropinone-derived alkaloids as potent anticancer agents: Synthesis, tyrosinase inhibition, mechanism of action, DFT calculation, and molecular docking studies. Int. J. Mol. Sci. 21(23), 9050 (2020).
- 3. Majewski, M. and Lazny, R. Synthesis of tropane alkaloids via enantioselective deprotonation of tropinone. J. Org. Chem. 60(18), 5825-5830 (1995).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 07/01/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM