



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)


# PRODUCT INFORMATION



## 7-Methylxanthine

Item No. 34348

**CAS Registry No.:** 552-62-5  
**Formal Name:** 3,7-dihydro-7-methyl-1H-purine-2,6-dione  
**Synonyms:** 7-MX, NSC 7861  
**MF:** C<sub>6</sub>H<sub>6</sub>N<sub>4</sub>O<sub>2</sub>  
**FW:** 166.1  
**Purity:** ≥98%  
**Supplied as:** A crystalline solid  
**Storage:** -20°C  
**Stability:** ≥4 years  
**Item Origin:** Synthetic



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

7-Methylxanthine is supplied as a crystalline solid. A stock solution may be made by dissolving the 7-methylxanthine in the solvent of choice, which should be purged with an inert gas. 7-Methylxanthine is slightly soluble in ethanol and DMSO.

### Description

7-Methylxanthine is an adenosine A<sub>1</sub> receptor antagonist and active metabolite of theobromine (Item No. 21745) and caffeine (Item No. 14118).<sup>1,2</sup> It inhibits binding of the adenosine A<sub>1</sub> receptor agonist cyclohexyladenosine in rat cerebral cortical membranes in a concentration-dependent manner.<sup>2</sup> 7-Methylxanthine (300 mg/kg) inhibits eye elongation and reduces decreases in the diameter of scleral collagen fibrils in a guinea pig model of form-deprivation myopia.<sup>3</sup>

### References

1. Cornish, H.H. and Christman, A.A. A study of the metabolism of theobromine, theophylline, and caffeine in man. *J. Biol. Chem.* **228**(1), 315-323 (1957).
2. Daly, J.W., Butts-Lamb, P., and Padgett, W. Subclasses of adenosine receptors in the central nervous system: Interaction with caffeine and related methylxanthines. *Cell. Mol. Neurobiol.* **3**(1), 69-80 (1983).
3. Cui, D., Trier, K., Zeng, J., et al. Effects of 7-methylxanthine on the sclera in form deprivation myopia in guinea pigs. *Acta. Ophthalmol.* **89**(4), 328-334 (2011).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/04/2022

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM