

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Corylifol A

Item No. 34616

CAS Registry No.: 775351-88-7

Formal Name: 3-[3-[(2E)-3,7-dimethyl-2,6-octadien-1-

yl]-4-hydroxyphenyl]-7-hydroxy-4H-1-

benzopyran-4-one

Synonym: Corylinin $C_{25}H_{26}O_4$ MF: FW: 390.5 **Purity:** ≥98% λ_{max} : 249 nm UV/Vis.: Supplied as: A solid -20°C Storage: Stability: ≥2 years

Plant/Psoralea corylifolia Item Origin:

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Corylifol A is supplied as a solid. A stock solution may be made by dissolving the corylifol A in the solvent of choice, which should be purged with an inert gas. Corylifol A is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of corylifol A in ethanol is approximately 25 mg/ml and approximately 30 mg/ml in DMSO and DMF.

Corylifol A is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, corylifol A should first be dissolved in DMSO and then diluted with the agueous buffer of choice. Corylifol A has a solubility of approximately 0.30 mg/ml in a 1:2 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Corylifol A is a flavonoid that has been found in P. corylifolia and has diverse biological activities. 1-4 It inhibits the activity of carboxylesterase 2 (CES2; $K_i = 0.62 \mu M$ for the human enzyme), as well as induces estrogen receptor α (ER α) and ER β transcriptional activation in reporter assays using HeLa cells when used at concentrations ranging from 0.1 to 10 μ M.^{2,3} Corylifol A is cytotoxic to (IC₅₀ = 13.5 μ g/ml), and inhibits IL-6-induced STAT3 promoter activity in (IC₅₀ = 0.81 μ M in a reporter assay), Hep3B hepatocellular carcinoma cells.^{1,4}

References

- 1. Lee, S.W., Yun, B.R., Kim, M.H., et al. Phenolic compounds isolated from Psoralea corylifolia inhibit IL-6-induced STAT3 activation. Planta Med. 78(9), 903-906 (2012).
- 2. Li, Y.-G., Hou, J., Li, S.-Y., et al. Fructus Psoraleae contains natural compounds with potent inhibitory effects towards human carboxylesterase 2. Fitoterapia 101, 99-106 (2015).
- Xin, D., Wang, H., Yang, J., et al. Phytoestrogens from Psoralea corylifolia reveal estrogen receptor-subtype selectivity. Phytomedicine 17(2), 126-131 (2010).
- 4. Song, P., Yang, X.-Z., and Yuan, J.-Q. Cytotoxic constituents from Psoralea corylifolia. J. Asian Nat. Prod. Res. **15(6)**, 624-630 (2013).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the mater can be found on our website.

Copyright Cayman Chemical Company, 08/18/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM