

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Sophoraflavanone G

Item No. 34755

CAS Registry No.:	97938-30-2
Formal Name:	(2S)-2-(2,4-dihydroxyphenyl)-2,3-
	dihydro-5,7-dihydroxy-8-[(2R)-5-
	methyl-2-(1-methylethenyl)-4-hexen-
	1-yl]-4H-1-benzopyran-4-one
Synonyms:	Kushenol F, Norkurarinone, Vexibinol
MF:	$C_{25}H_{28}O_6$
FW:	424.5
Purity:	≥95% Он
Supplied as:	A solid
Storage:	-20°C Ö
Stability:	≥2 years
Item Origin:	Plant/Sophora flavescens
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.	

Laboratory Procedures

Sophoraflavanone G is supplied as a solid. A stock solution may be made by dissolving the sophoraflavanone G in the solvent of choice, which should be purged with an inert gas. Sophoraflavanone G is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of sophoraflavanone G in ethanol is approximately 20 mg/ml and approximately 30 mg/ml in DMSO and DMF.

Description

Sophoraflavanone G is a flavonoid that has been found in S. flavescens and has diverse biological activities.¹⁻⁴ It is active against 21 strains of methicillin-resistant S. aureus (MRSA; MICs = 3.13-6.25 μ g/ml) and inhibits the activities of α -glucosidase and β -amylase (K_is = 5.6 and 30.6 μ M, respectively).^{1,2} Sophoraflavanone G is cytotoxic to HL-60 and HepG2 cancer cells (IC₅₀s = 12.5 and 13.3 μ M, respectively) and induces apoptosis in HL-60 cells when used at concentrations of 10 or 25 μ M.³ It reduces LPS-induced production of prostaglandin E₂ (PGE₂; Item No. 14010) in RAW 264.7 cells when used at concentrations ranging from 1 to 50 μ M.⁴ Sophoraflavanone G (250 mg/kg) inhibits carrageenan-induced paw edema in rats.

References

- 1. Sato, M., Tsuchiya, H., Takase, I., et al. Antibacterial activity of flavanone isolated from Sophora exigua against methicillin-resistant Staphylococcus aureus and its combination with antibiotics. Phytother. Res. 9(7), 509-512 (1995).
- 2. Kim, J.H., Ryu, Y.B., Kang, N.S., et al. Glycosidase inhibitory flavonoids from Sophora flavescens. Biol. Pharm. Bull. 29(2), 302-305 (2006).
- 3. Ko, W.G., Kang, T.H., Kim, N.Y., et al. Lavandulylfavonoids: A new class of in vitro apoptogenic agents from Sophora favescens. Toxicol. In Vitro 14(5), 429-433 (2000).
- Kim, D.W., Chi, Y.S., Son, K.H., et al. Effects of sophoraflavanone G, a prenylated flavonoid from Sophora 4. flavescens, on cyclooxygenase-2 and in vivo inflammatory response. Arch. Pharm. Res. 25(3), 329-335 (2002).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 08/05/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM