

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



3-Hydroxycoumarin

Item No. 34815

CAS Registry No.: 939-19-5

Formal Name: 3-hydroxy-2H-1-benzopyran-2-one

Synonyms: 3-Coumarinol, NSC 74691

MF: $C_9H_6O_3$ FW: 162.1 **Purity:** ≥95%

UV/Vis.: λ_{max} : 230, 309 nm

Supplied as: A solid Storage: -20°C Stability: ≥4 years Item Origin: Synthetic

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

3-Hydroxycoumarin is supplied as a solid. A stock solution may be made by dissolving the 3-hydroxycoumarin in the solvent of choice, which should be purged with an inert gas, 3-Hydroxycoumarin is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of 3-hydroxycoumarin in these solvents is approximately 1, 10, and 15 mg/ml, respectively.

3-Hydroxycoumarin is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, 3-hydroxycoumarin should first be dissolved in DMF and then diluted with the aqueous buffer of choice. 3-Hydroxycoumarin has a solubility of approximately 0.20 mg/ml in a 1:4 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

3-Hydroxycoumarin is a coumarin that has been found in A. reinwardtii and has diverse biological activities. $^{1-3}$ It inhibits 15-lipoxygenase-1 (15-LO-1; IC $_{50}$ = 9.5 μ M). 1 3-hydroxycoumarin scavenges DPPH (Item No. 14805) radicals in a cell-free assay ($IC_{50} = 0.61$ mM) and inhibits lipid peroxidation in vitro (IC₅₀ = 69.07 mM).² It is protective against UVB-induced embryotoxicity in sea urchins (E. lucunter).³

References

- 1. Alavi, S.J., Sadeghian, H., Seyedi, S.M., et al. A novel class of human 15-LOX-1 inhibitors based on 3-hydroxycoumarin. Chem. Biol. Drug Des. 91(6), 1125-1132 (2017).
- 2. Rattanapan, J., Sichaem, J., and Tip-pyang, S. Chemical constituents and antioxidant activity from the stems of Alyxia reinwardtii. Rec. Nat. Prod. 6(3), 288-291 (2012).
- de Araujo Leite, J.C., de Castro, T.M.X., Barbosa-Filho, J.M., et al. Photoprotective effect of coumarin and 3-hydroxycoumarin in sea urchin gametes and embryonic cells. J. Photochem. Photobiol. B Biol. 146, 44-51 (2015).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the m can be found on our website.

Copyright Cayman Chemical Company, 12/08/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM