

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



N⁴-Acetylcytidine

Item No. 35219

CAS Registry No.: 3768-18-1 Formal Name: N-acetyl-cytidine

ac⁴C Synonym:

MF: $C_{11}H_{15}N_3O_6$ FW: 285.3 **Purity:**

 λ_{max} : 215, 247, 299 nm UV/Vis.:

A solid Supplied as: -20°C Storage: Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

N⁴-Acetylcytidine is supplied as a solid. A stock solution may be made by dissolving the N⁴-acetylcytidine in the solvent of choice, which should be purged with an inert gas. N^4 -Acetylcytidine is soluble in the organic solvent DMSO at a concentration of approximately 1 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of N⁴-acetylcytidine can be prepared by directly dissolving the solid in aqueous buffers. The solubility of N⁴-acetylcytidine in PBS (pH 7.2) is approximately 2 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

N⁴-Acetylcytidine is a catabolite of cytidine.¹ It activates BV-2 microglia when used at a concentration of 0.3 mM, an effect that can be blocked by the adenosine A_{2A} receptor antagonist SCH 58261 (Item No. 19676).2 It also increases protein levels of the NOD-like receptor protein 3 (NLRP3) inflammasome, an effect that can be blocked by high mobility group box 1 (HMGB1) siRNA in BV-2 microglia. Urine levels of N⁴-acetylcytidine are increased in mice with tumors induced by 3-methylcholanthrene. 1 N⁴-Acetylcytidine is also found as a post-transcriptional modification in RNA.3

References

- 1. Thomale, J. and Nass, G. Elevated urinary excretion of RNA catabolites as an early signal of tumor development in mice. Cancer Lett. 15(2), 149-159 (1982).
- Duan, J.J., Zhang, Q., Hu, X., et al. N⁴-acetylcytidine is required for sustained NLRP3 inflammasome activation via HMGB1 pathway in microglia. Cell. Signal. 58, 44-52 (2019).
- Bartee, D., Nance, K.D., and Meier, J.L. Site-specific synthesis of N⁴-acetylcytidine in RNA reveals physiological duplex stabilization. J. Am. Chem. Soc. 144(8), 3487-3496 (2022).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 12/12/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM