

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Monoacetylcadaverine (hydrochloride)

Item No. 35275

CAS Registry No.: Formal Name:	102029-76-5 N-(5-aminopentyl)-acetamide, monohydrochloride	0
MF:	$C_7H_{16}N_2O \bullet HCI$	Ŭ ^ ^ ^
FW:	180.7	N NH ₂
Purity:	≥95%	H • HCI
Supplied as:	A solid	
Storage:	-20°C	
Stability:	≥4 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.		

Laboratory Procedures

Monoacetylcadaverine (hydrochloride) is supplied as a solid. A stock solution may be made by dissolving the monoacetylcadaverine (hydrochloride) in the solvent of choice, which should be purged with an inert gas. Monoacetylcadaverine (hydrochloride) is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of monoacetylcadaverine (hydrochloride) in these solvents is approximately 12, 16, and 14 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of monoacetylcadaverine (hydrochloride) can be prepared by directly dissolving the solid in aqueous buffers. The solubility of monoacetylcadaverine (hydrochloride) in PBS (pH 7.2) is approximately 3 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Monoacetylcadaverine is a metabolite of the endogenous polyamine cadaverine.¹ It is formed from cadaverine via acetylation in the nuclear and microsomal fractions of isolated mouse brain.² Plasma levels of monoacetylcadaverine are increased in patients with schizophrenia.³

References

- 1. van den Berg, G.A., Muskiet, F.A., Kingma, A.W., et al. Simultaneous gas-chromatographic determination of free and acetyl-conjugated polyamines in urine. Clin. Chem. 32(10), 1930-1937 (1986).
- 2. Ortiz, J.G., Giacobini, E., and Schmidt-Glenewinkel, T. Acetylation of polyamines in mouse brain: Subcellular and regional distribution. J. Neurosci. Res. 9(2), 193-201 (1983).
- 3. Dolezalova, H. and Kauco, M.S. Monoacylcadaverines in the blood of schizophrenic patients. J. Chromatogr. 146(1), 67-76 (1978).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/12/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM