

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



N⁶,N⁶-Dimethyladenosine

Item No. 35348

CAS Registry No.: 2620-62-4

6-N,N-Dimethyladenosine, Synonyms:

6-DMA, m⁶₂A

MF: $C_{12}H_{17}N_5O_4$ 295.3 FW: **Purity:**

UV/Vis.: λ_{max}: 216, 274 nm

A solid Supplied as: -20°C Storage: Stability: ≥4 years Item Origin: Synthetic

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

N⁶,N⁶-Dimethyladenosine is supplied as a solid. A stock solution may be made by dissolving the N⁶,N⁶-dimethyladenosine in the solvent of choice, which should be purged with an inert gas. N⁶,N⁶-Dimethyladenosine is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of N⁶,N⁶-dimethyladenosine in these solvents is approximately 25 and 20 mg/ml, respectively. N⁶,N⁶-Dimethyladenosine is slightly soluble in ethanol.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of N⁶,N⁶-dimethyladenosine can be prepared by directly dissolving the solid in aqueous buffers. The solubility of N⁶,N⁶-dimethyladenosine in PBS (pH 7.2) is approximately 2 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

N⁶,N⁶-Dimethyladenosine is a modified nucleoside.¹ It has been found in tRNA isolated from M. bovis. Dimethylation of adjacent adenosine nucleosides (m⁶₂Am⁶₂A) is found in E. coli 16S rRNA and is important for the initiation of protein synthesis. ² N⁶, N⁶-Dimethyladenosine inhibits the proliferation of L1210 leukemia cells in vitro ($IC_{50} = 0.5 \mu g/mI$) but does not have antitumor activity in vivo.² It is also a component of the antibiotic and protein synthesis inhibitor puromycin (Item No. 13884).

References

- 1. Chan, C.T.Y., Chionh, Y.H., Ho, C.-H., et al. Identification of N⁶,N⁶-dimethyladenosine in transfer RNA from Mycobacterium bovis Bacille Calmette-Guérin. Molecules 16(6), 5168-5181 (2011).
- 2. Poldermans, B., Van Buul, C.P.J.J., and Van Knippenberg, P.H. Studies on the function of two adjacent N⁶,N⁶-dimethyladenosines near the 3' end of 16 S ribosomal RNA of Escherichia coli. II. The effect of the absence of the methyl groups on initiation of protein biosynthesis. J. Biol. Chem. 254(18), 9090-9093 (1979).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 12/12/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM